

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 502 589 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
26.09.2001 Bulletin 2001/39

(51) Int Cl.7: **C12Q 1/68, C12P 19/34**
// C07H21/04

(21) Application number: **92201245.5**

(22) Date of filing: **27.03.1986**

(54) Kit for use in amplifying and detecting nucleic acid sequences

Kit zur Verwendung bei der Amplifizierung und beim Nachweis von Nukleinsäuresequenzen

Trousse pour utilisation dans l'amplification et la détection de séquences d'acide nucleique

(84) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(30) Priority: **28.03.1985 US 716975**
25.10.1985 US 791308
07.02.1986 US 828144

(43) Date of publication of application:
09.09.1992 Bulletin 1992/37

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
86302298.4 / O 200 362

(73) Proprietor: **F. HOFFMANN-LA ROCHE AG**
4002 Basel (CH)

(72) Inventors:
• **Mullis, Kary Banks**
Kensington, California 94708 (US)
• **Arnheim, Norman**
Woodland Hills, California 91364 (US)
• **Saiki, Randall Keichi**
Rochmond, California 94805 (US)

• **Erlich, Henry Anthony**
Oakland, California 94602 (US)
• **Horn, Glenn Thomas**
Marlborough, Massachussets 07152 (US)
• **Scharf, Stephen Joel**
Berkeley, California 94709 (US)

(74) Representative: **Jaenichen, Hans-Rainer, Dr. et al**
Vossius & Partner,
Postfach 86 07 67
81634 München (DE)

(56) References cited:
EP-A- 0 086 548 EP-A- 0 138 242
EP-A- 0 139 501 EP-A- 0 155 188
US-A- 4 351 901

• **JOURNAL MOLECULAR BIOOLOGY vol. 56 , 28**
March 1971 , US pages 341 - 361 KLEPPE, K. ET
AL 'repair replication of short synthetic DNA's
as catalysed by DNA Polymerases'
• **JOURNAL OF BIOLOGICAL CHEMISTRY. vol.**
249 , August 1974 , BALTIMORE US pages 5213
- 5221 PANET A ET AL 'studies on
polynucleotides'

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 502 589 B1

Description

[0001] The present invention relates to a kit for use in amplifying existing nucleic acid sequences if they are present in the test sample and detecting them if present by using a probe. The invention can be used by employing a process for producing any particular nucleic acid sequence from a given sequence of DNA or RNA in amounts which are large compared to the amount initially present so as to facilitate detection of the sequences. The DNA or RNA may be single- or double-stranded, and may be a relatively pure species or a component of a mixture of nucleic acids. The process utilizes a repetitive reaction to accomplish the amplification of the desired nucleic acid sequence.

[0002] For diagnostic applications in particular, the target nucleic acid sequence may be only a small portion of the DNA or RNA in question, so that it may be difficult to detect its presence using nonisotopically labeled or end-labeled oligonucleotide probes. Much effort is being expended in increasing the sensitivity of the probe detection systems, but little research has been conducted on amplifying the target sequence so that it is present in quantities sufficient to be readily detectable using currently available methods.

[0003] Several methods have been described in the literature for the synthesis of nucleic acids de novo or from an existing sequence. These methods are capable of producing large amounts of a given nucleic acid of completely specified sequence.

[0004] One known method for synthesizing nucleic acids de novo involves the organic synthesis of a nucleic acid from nucleoside derivatives. This synthesis may be performed in solution or on a solid support. One type of organic synthesis is the phosphotriester method, which has been utilized to prepare gene fragments or short genes. In the phosphotriester method, oligonucleotides are prepared which can then be joined together to form longer nucleic acids. For a description of this method, see Narang, S.A., et al., *Meth. Enzymol.*, 68, 90 (1979) and U.S. Patent No. 4,356,270. The patent describes the synthesis and cloning of the somatostatin gene.

[0005] A second type of organic synthesis is the phosphodiester method, which has been utilized to prepare a tRNA gene. See Brown, E.L., et al., *Meth. Enzymol.*, 68, 109 (1979) for a description of this method. As in the phosphotriester method, the phosphodiester method involves synthesis of oligonucleotides which are subsequently joined together to form the desired nucleic acid.

[0006] Although the above processes for de novo synthesis may be utilized to synthesize long strands of nucleic acid, they are not very practical to use for the synthesis of large amounts of a nucleic acid. Both processes are laborious and time-consuming, require expensive equipment and reagents, and have a low overall efficiency. The low overall efficiency may be caused by the inefficiencies of the synthesis of the oligonucleotides and of the joining reactions. In the synthesis of a long nucleic acid, or even in the synthesis of a large amount of a shorter nucleic acid, many oligonucleotides would need to be synthesized and many joining reactions would be required. Consequently, these methods would not be practical for synthesizing large amounts of any desired nucleic acid.

[0007] Methods also exist for producing nucleic acids in large amounts from small amounts of the initial existing nucleic acid. These methods involve the cloning of a nucleic acid in the appropriate host system, where the desired nucleic acid is inserted into an appropriate vector which is used to transform the host. When the host is cultured the vector is replicated, and hence more copies of the desired nucleic acid are produced. For a brief description of sub-cloning nucleic acid fragments, see Maniatis, T., et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, pp. 390-401 (1982). See also the techniques described in U.S. Patent Nos. 4,416,988 and 4,403,036.

[0008] A third method for synthesizing nucleic acids, described in U.S. Patent No. 4,293,652, is a hybrid of the above-described organic synthesis and molecular cloning methods. In this process, the appropriate number of oligonucleotides to make up the desired nucleic acid sequence is organically synthesized and inserted sequentially into a vector which is amplified by growth prior to each succeeding insertion.

[0009] The process which can be carried out by using the kit of the present invention bears some similarity to the molecular cloning method; however, it does not involve the propagation of any organism and thereby avoids the possible hazards or inconvenience which this entails. The process also does not require synthesis of nucleic acid sequences unrelated to the desired sequence, and thereby obviates the need for extensive purification of the product from a complicated biological mixture.

[0010] In *J.Mol.Biol.*, 56 (1971), 341-361, Kleppe et al discuss primer extension reactions using templates corresponding to portions of a tRNA gene, in which reactions the primers are used are complementary to substantial parts of corresponding templates and are extended therealong thereby to provide duplex DNAs. These template copying reactions, involving simple primer extension, are termed "repair replication" by the authors. The final paragraph of the article theorises that if duplex DNA denaturation is effected in the presence of appropriate primers, two structures consisting of a full length template strand complexed to a primer could be produced upon cooling, and repair replication achieved by adding DNA polymerase. - The paragraph suggests that this process could be repeated.

[0011] However, there is no detailed explanation of the precise techniques to be carried out, nor any discussion of which primers are "appropriate", and the possibility of a problem of template renaturation (reforming a duplex) is discussed with the suggestion that, if necessary, strand separation would have to be resorted to with subsequent repair

replication.

[0012] The kit of the present invention is used in process for amplifying one or more specific nucleic acid sequences present in a nucleic acid or mixture thereof using primers and agents for polymerization and then detecting the amplified sequence. The extension product of one primer when hybridized to the other becomes a template for the production of the desired specific nucleic acid sequence, and vice versa, and the process is repeated as often as is necessary to produce the desired amount of the sequence. This method is expected to be more efficient than the methods described above for producing large amounts of nucleic acid from a target sequence and to produce such nucleic acid in a comparatively short period of time. The method is especially useful for amplifying rare species of nucleic acid present in a mixture of nucleic acids for effective detection of such species.

[0013] More specifically, the present invention provides an exponential amplification and detection kit for the amplification and detection of (a) specific template nucleic acid sequence(s) contained in a single- or double-stranded nucleic acid or in a mixture of such nucleic acids in a sample, which kit comprises in packaged form:

(a) at least a first and a second oligonucleotide primer different from each other, wherein

(aa) one of said primers is substantially complementary to said single-stranded nucleic acid or to one strand of said double-stranded nucleic acid,

(ab) the other primer of said primers is substantially complementary to a complement of said single-stranded nucleic acid or to the other strand of said double-stranded nucleic acid, and wherein

(ac) said primers define the termini of the specific nucleic acid sequence to be amplified and detected;

(b) an agent for polymerization; and

(c) means for detecting the amplified specific nucleic acid sequence.

[0014] Particular embodiments of the above kit are defined in claims 2 to 12.

[0015] The invention further provides the use of a kit as above for enabling detection and/or characterization of specific nucleic acid sequences associated with infectious diseases such as those caused by bacteria, viruses and protozoan parasites, genetic disorders such as those caused by specific deletions and/or mutations in genomic DNA or cellular disorders such as cancer.

[0016] Co-pending European Application No 201184 discloses processes for amplifying nucleic acid sequences, and this Application is divided from European Application No 86 302298.4.

[0017] Figure 1 illustrates a 94 base pair length sequence of human β -globin desired to be amplified. The single base pair change which is associated with sickle cell anemia is depicted beneath the 94-mer.

[0018] Figure 2 illustrates a photograph of an ethidium bromide-stained polyacrylamide gel demonstrating amplification of the 94-mer contained in human wild-type DNA and in a plasmid containing a 1.9 kb BamHI fragment of the normal β -globin gene (designated pBR328:HbA).

[0019] Figure 3 illustrates a photograph of an ethidium bromide-stained polyacrylamide gel demonstrating amplification of any of the specific target 94-mer sequence present in pBR328:HbA, a plasmid containing a 1.9 kb BamHI fragment of the sickle cell allele of β -globin (designated pBR328:HbS), pBR328:HbA where the sequence to be amplified is cleaved with MstII, and pBR328:HbS where the sequence to be amplified has been treated but not cleaved with MstII.

[0020] Figure 4 illustrates in detail the steps and products of the polymerase chain reaction for amplification of the desired 94-mer sequence of human β -globin for three cycles using two oligonucleotide primers.

[0021] Figure 5 represents a photograph of an ethidium bromide-stained polyacrylamide gel demonstrating amplification after four cycles of a 240-mer sequence in pBR328:HbA, where the aliquots are digested with NcoI (Lane 3), MstII (Lane 4) or HinFI (Lane 5). Lane 1 is the molecular weight standard and Lane 2 contains the intact 240-bp product.

[0022] Figure 6 illustrates the sequence of the normal (β^A) and sickle cell (β^S) β -globin genes in the region of the DdeI and HinFI restriction sites, where the single lines for β^A mark the position of the DdeI site (CTGAG) and the double bars for β^A and β^S mark the position of the HinFI site (GACTC).

[0023] Figure 7 illustrates the results of sequential digestion of normal β -globin using a 40-mer probe and DdeI followed by HinFI restriction enzymes.

[0024] Figure 8 illustrates the results of sequential digestion of sickle β -globin using the same 40-mer probe as in Figure 7 and DdeI followed by HinFI restriction enzymes.

[0025] Figure 9 illustrates a photograph of an ethidium bromide-stained polyacrylamide gel demonstrating the use of the same 40-mer probe as in Figure 7 to specifically characterize the beta-globin alleles present in samples of whole human DNA which have been subjected to amplification, hybridization with the probe, and sequential digestion with DdeI and HinFI.

[0026] Figure 10 illustrates a photograph of a 6% NuSieve agarose gel visualized using ethidium bromide and UV

light. This photograph demonstrates amplification of a sub-fragment of a 110-bp amplification product which sub-fragment is an inner nested set within the 110-bp fragment.

[0027] The term "oligonucleotide" as used herein in referring to primers, probes, oligomer fragments to be detected, oligomer controls and unlabeled blocking oligomers is defined as a molecule comprised of two or more deoxyribonucleotides or ribonucleotides, preferably more than three. Its exact size will depend on many factors, which in turn depend on the ultimate function or use of the oligonucleotide.

[0028] The term "primer" as used herein refers to an oligonucleotide whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product which is complementary to a nucleic acid strand is induced, i.e., in the presence of nucleotides and an agent for polymerization such as DNA polymerase and at a suitable temperature and pH. The primer is preferably single stranded for maximum efficiency in amplification, but may alternatively be double stranded. If double stranded, the primer is first treated to separate its strands before being used to prepare extension products. Preferably, the primer is an oligodeoxyribonucleotide. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the agent for polymerization. The exact lengths of the primers will depend on many factors, including temperature and source of primer. For example, depending on the complexity of the target sequence, the oligonucleotide primer typically contains 15-25 or more nucleotides, although it may contain fewer nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with template.

[0029] The primers herein are selected to be "substantially" complementary to the different strands of each specific sequence to be amplified. This means that the primers must be sufficiently complementary to hybridize with their respective strands. Therefore, the primer sequence need not reflect the exact sequence of the template. For example, a non-complementary nucleotide fragment may be attached to the 5' end of the primer, with the remainder of the primer sequence being complementary to the strand. Alternatively, non-complementary bases or longer sequences can be interspersed into the primer, provided that the primer sequence has sufficient complementarity with the sequence of the strand to be amplified to hybridize therewith and thereby form a template for synthesis of the extension product of the other primer.

[0030] As used herein, the terms "restriction endonucleases" and "restriction enzymes" refer to bacterial enzymes each of which cut double-stranded DNA at or near a specific nucleotide sequence.

[0031] As used herein, the term "DNA polymorphism" refers to the condition in which two or more different nucleotide sequences can exist at a particular site in DNA.

[0032] The term "restriction fragment length polymorphism" ("RFLP") refers to the differences among individuals in the lengths of restriction fragments formed by digestion with a particular restriction endonuclease.

[0033] The present invention is directed to a kit for amplifying and detecting any one or more desired specific nucleic acid sequences suspected of being in a nucleic acid. Because large amounts of a specific sequence may be produced by this kit, the present invention may be used for improving the efficiency of cloning DNA or messenger RNA and for amplifying a target sequence to facilitate detection thereof.

[0034] In general, the process carried out by the present kit involves a chain reaction for producing, in exponential quantities relative to the number of reaction steps involved, at least one specific nucleic acid sequence given (a) that the ends of the required sequence are known in sufficient detail that oligonucleotides can be synthesized which will hybridize to them, and (b) that a small amount of the sequence is available to initiate the chain reaction. The product of the chain reaction will be a discrete nucleic acid duplex with termini corresponding to the ends of the specific primers employed.

[0035] Any source of nucleic acid, in purified or nonpurified form, can be utilized as the starting nucleic acid or acids, provided it is suspected of containing the specific nucleic acid sequence desired. Thus, the process may employ, for example, DNA or RNA, including messenger RNA, which DNA or RNA may be single stranded or double stranded. In addition, a DNA-RNA hybrid which contains one strand of each may be utilized. A mixture of any of these nucleic acids may also be employed, or the nucleic acids produced from a previous amplification reaction herein using the same or different primers may be so utilized. The specific nucleic acid sequence to be amplified may be only a fraction of a larger molecule or can be present initially as a discrete molecule, so that the specific sequence constitutes the entire nucleic acid. It is not necessary that the sequence to be amplified be present initially in a pure form; it may be a minor fraction of a complex mixture, such as a portion of the β -globin gene contained in whole human DNA or a portion of nucleic acid sequence due to a particular microorganism which organism might constitute only a very minor fraction of a particular biological sample. The starting nucleic acid may contain more than one desired specific nucleic acid sequence which may be the same or different. Therefore, the present process is useful not only for producing large amounts of one specific nucleic acid sequence, but also for amplifying simultaneously more than one different specific nucleic acid sequence located on the same or different nucleic acid molecules.

[0036] The nucleic acid or acids may be obtained from any source, for example, from plasmids such as pBR322, from cloned DNA or RNA, or from natural DNA or RNA from any source, including bacteria, yeast, viruses, and higher

organisms such as plants or animals. DNA or RNA may be extracted from blood, tissue material such as chorionic villi or amniotic cells by a variety of techniques such as that described by Maniatis et al., Molecular Cloning (1982), 280-281.

[0037] Any specific nucleic acid sequence can be produced by the present kit. It is only necessary that a sufficient number of bases at both ends of the sequence be known in sufficient detail so that two oligonucleotide primers can be prepared which will hybridize to different strands of the desired sequence and at relative positions along the sequence such that an extension product synthesized from one primer, when it is separated from its template (complement), can serve as a template for extension of the other primer into a nucleic acid of defined length. The greater the knowledge about the bases at both ends of the sequence, the greater can be the specificity of the primers for the target nucleic acid sequence, and thus the greater the efficiency of the process. It will be understood that the word primer as used hereinafter may refer to more than one primer, particularly in the case where there is some ambiguity in the information regarding the terminal sequence(s) of the fragment to be amplified. For instance, in the case where a nucleic acid sequence is inferred from protein sequence information a collection of primers containing sequences representing all possible codon variations based on degeneracy of the genetic code will be used for each strand. One primer from this collection will be homologous with the end of the desired sequence to be amplified.

[0038] The oligonucleotide primers may be prepared using any suitable method, such as, for example, the phosphotriester and phosphodiester methods described above, or automated embodiments thereof. In one such automated embodiment diethylphosphoramidites are used as starting materials and may be synthesized as described by Beaucage et al., Tetrahedron Letters (1981), 22:1859-1862. One method for synthesizing oligonucleotides on a modified solid support is described in U.S. Patent No. 4,458,066. It is also possible to use a primer which has been isolated from a biological source (such as a restriction endonuclease digest).

[0039] The specific nucleic acid sequence is produced by using the nucleic acid containing that sequence as a template. If the nucleic acid contains two strands, it is necessary to separate the strands of the nucleic acid before it can be used as the template, either as a separate step or simultaneously with the synthesis of the primer extension products. This strand separation can be accomplished by any suitable denaturing method including physical, chemical or enzymatic means. One physical method of separating the strands of the nucleic acid involves heating the nucleic acid until it is completely (>99%) denatured. Typical heat denaturation may involve temperatures ranging from about 80 to 105°C for times ranging from about 1 to 10 minutes. Strand separation may also be induced by an enzyme from the class of enzymes known as helicases or the enzyme RecA, which has helicase activity and in the presence of riboATP is known to denature DNA. The reaction conditions suitable for separating the strands of nucleic acids with helicases are described by Kuhn Hoffmann-Berling, CSH-Quantitative Biology, 43:63 (1978), and techniques for using RecA are reviewed in C. Radding, Ann. Rev. Genetics, 16:405-37 (1982).

[0040] If the original nucleic acid containing the sequence to be amplified is single stranded, its complement is synthesized by adding one or two oligonucleotide primers thereto. If an appropriate single primer is added, a primer extension product is synthesized in the presence of the primer, an agent for polymerization and the four nucleotides described below. The product will be partially complementary to the single-stranded nucleic acid and will hybridize with the nucleic acid strand to form a duplex of unequal length strands that may then be separated into single strands as described above to produce two single separated complementary strands. Alternatively, two appropriate primers may be added to the single-stranded nucleic acid and the reaction carried out.

[0041] If the original nucleic acid constitutes the sequence to be amplified, the primer extension product(s) produced will be completely complementary to the strands of the original nucleic acid and will hybridize therewith to form a duplex of equal length strands to be separated into single-stranded molecules.

[0042] When the complementary strands of the nucleic acid or acids are separated, whether the nucleic acid was originally double or single stranded, the strands are ready to be used as a template for the synthesis of additional nucleic acid strands. This synthesis can be performed using any suitable method. Generally it occurs in a buffered aqueous solution, preferably at a pH of 7-9, most preferably about 8. Preferably, a molar excess (for cloned nucleic acid, usually about 1000:1 primer:template, and for genomic nucleic acid, usually about 10⁶:1 primer:template) of the two oligonucleotide primers is added to the buffer containing the separated template strands. It is understood, however, that the amount of complementary strand may not be known if the kit herein is used for diagnostic applications, so that the amount of primer relative to the amount of complementary strand cannot be determined with certainty. As a practical matter, however, the amount of primer added will generally be in molar excess over the amount of complementary strand (template) when the sequence to be amplified is contained in a mixture of complicated long-chain nucleic acid strands. A large molar excess is preferred to improve the efficiency of the process.

[0043] The deoxyribonucleoside triphosphates dATP, dCTP, dGTP and TTP are also added to the synthesis mixture in adequate amounts and the resulting solution is heated to about 90-100°C for from about 1 to 10 minutes, preferably from 1 to 4 minutes. After this heating period the solution is allowed to cool to from 20-40°C, which is preferable for the primer hybridization. To the cooled mixture is added an agent for polymerization, and the reaction is allowed to occur under conditions known in the art. This synthesis reaction may occur at from room temperature up to a temperature above which the agent for polymerization no longer functions efficiently. Thus, for example, if DNA polymerase

is used as the agent for polymerization, the temperature is generally no greater than about 45°C. Preferably an amount of dimethylsulfoxide (DMSO) is present which is effective in detection of the signal or the temperature is 35-40°C. Most preferably, 5-10% by volume DMSO is present and the temperature is 35-40°C. For certain applications, where the sequences to be amplified are over 110 base pair fragments, such as the HLA DQ- α or - β genes, an effective amount (e.g., 10% by volume) of DMSO is added to the amplification mixture, and the reaction is carried at 35-40°C, to obtain detectable results or to enable cloning.

[0044] The agent for polymerization may be any compound or system which will function to accomplish the synthesis of primer extension products, including enzymes. Suitable enzymes for this purpose include, for example, *E. coli* DNA polymerase I, Klenow fragment of *E. coli* DNA polymerase I, T4 DNA polymerase, other available DNA polymerases, reverse transcriptase, and other enzymes, including heat-stable enzymes, which will facilitate combination of the nucleotides in the proper manner to form the primer extension products which are complementary to each nucleic acid strand. Generally, the synthesis will be initiated at the 3' end of each primer and proceed in the 5' direction along the template strand, until synthesis terminates, producing molecules of different lengths. There may be agents, however, which initiate synthesis at the 5' end and proceed in the other direction, using the same process as described above.

[0045] The newly synthesized strand and its complementary nucleic acid strand form a double-stranded molecule which is used in the succeeding steps of the process. In the next step, the strands of the double-stranded molecule are separated using any of the procedures described above to provide single-stranded molecules.

[0046] New nucleic acid is synthesized on the single-stranded molecules. Additional inducing agent, nucleotides and primers may be added if necessary for the reaction to proceed under the conditions prescribed above. Again, the synthesis will be initiated at one end of the oligonucleotide primers and will proceed along the single strands of the template to produce additional nucleic acid. After this step, half of the extension product will consist of the specific nucleic acid sequence bounded by the two primers.

[0047] The steps of strand separation and extension product synthesis can be repeated as often as needed to produce the desired quantity of the specific nucleic acid sequence. As will be described in further detail below, the amount of the specific nucleic acid sequence produced will accumulate in an exponential fashion.

[0048] When it is desired to produce more than one specific nucleic acid sequence from the first nucleic acid or mixture of nucleic acids, the appropriate number of different oligonucleotide primers are utilized. For example, if two different specific nucleic acid sequences are to be produced, four primers are utilized. Two of the primers are specific for one of the specific nucleic acid sequences and the other two primers are specific for the second specific nucleic acid sequence. In this manner, each of the two different specific sequences can be produced exponentially by the present process.

[0049] The present invention can be performed in a step-wise fashion where after each step new reagents are added, or simultaneously, where all reagents are added at the initial step, or partially step-wise and partially simultaneous, where fresh reagent is added after a given number of steps. If a method of strand separation, such as heat, is employed which will inactivate the agent for polymerization, as in the case of a heat-labile enzyme, then it is necessary to replenish the agent for polymerization after every strand separation step. The simultaneous method may be utilized when a number of purified components, including an enzymatic means such as helicase, is used for the strand separation step. In the simultaneous procedure, the reaction mixture may contain, in addition to the nucleic acid strand(s) containing the desired sequence, the strand-separating enzyme (e.g., helicase), an appropriate energy source for the strand-separating enzyme, such as rATP, the four nucleotides, the oligonucleotide primers in molar excess, and the inducing agent, e.g., Klenow fragment of *E. coli* DNA polymerase I. If heat is used for denaturation in a simultaneous process, a heat-stable inducing agent such as a thermostable polymerase may be employed which will operate at an elevated temperature, preferably 55-90°C depending on the inducing agent, at which temperature the nucleic acid will consist of single and double strands in equilibrium. For smaller lengths of nucleic acid, lower temperatures of about 50°C may be employed. The upper temperature will depend on the temperature at which the enzyme will degrade or the temperature above which an insufficient level of primer hybridization will occur. Such a heat-stable enzyme is described, e.g., by A. S. Kaledin et al., *Biokhimiya*, 45, 644-651 (1980). Each step of the process will occur sequentially notwithstanding the initial presence of all the reagents. Additional materials may be added as necessary. After the appropriate length of time has passed to produce the desired amount of the specific nucleic acid sequence, the reaction may be halted by inactivating the enzymes in any known manner or separating the components of the reaction.

[0050] The process carried out by use of the kit of the present invention may be conducted continuously. In one embodiment of an automated process, the reaction may be cycled through a denaturing region, a reagent addition region, and a reaction region. In another embodiment, the enzyme used for the synthesis of primer extension products can be immobilized in a column. The other reaction components can be continuously circulated by a pump through the column and a heating coil in series, thus the nucleic acids produced can be repeatedly denatured without inactivating the enzyme.

[0051] The process carried out by use of the kit of the present invention is demonstrated diagrammatically below where double-stranded DNA containing the desired sequence [S] comprised of complementary strands [S⁺] and [S⁻]

is utilized as the nucleic acid. During the first and each subsequent reaction cycle extension of each oligonucleotide primer on the original template will produce one new sDNA molecule product of indefinite length which terminates with only one of the primers. These products, hereafter referred to as "long products," will accumulate in a linear fashion; that is, the amount present after any number of cycles will be proportional to the number of cycles.

[0052] The long products thus produced will act as templates for one or the other of the oligonucleotide primers during subsequent cycles and will produce molecules of the desired sequence $[S^+]$ or $[S^-]$. These molecules will also function as templates for one or the other of the oligonucleotide primers, producing further $[S^+]$ and $[S^-]$, and thus a chain reaction can be sustained which will result in the accumulation of $[S]$ at an exponential rate relative to the number of cycles.

[0053] By-products formed by oligonucleotide hybridizations other than those intended are not self-catalytic (except in rare instances) and thus accumulate at a linear rate.

[0054] The specific sequence to be amplified, $[S]$, can be depicted diagrammatically as:

```

15      [S+]          5' AAAAAAAAAAXXXXXXXXXXCCCCCCCCC 3'
        [S-]          3' TTTTTTTTTTYYYYYYYYYYGGGGGGGGGG 5'

```

[0055] The appropriate oligonucleotide primers would be:

```

20      Primer 1:          GGGGGGGGGG
      Primer 2:          AAAAAA AAAA

```

so that if DNA containing $[S]$

```

30      ....ZZZZZZZZZZZZZZZZAAAAAAAAAAXXXXXXXXXXCCCCCCCCCZZZZZZZZZZZZZZZZ....
      ....ZZZZZZZZZZZZZZZZTTTTTTTTTTYYYYYYYYYYGGGGGGGGGGZZZZZZZZZZZZZZZZ....

```

is separated into single strands and its single strands are hybridized to Primers 1 and 2, the following extension reactions can be catalyzed by DNA polymerase in the presence of the four deoxyribonucleoside triphosphates:

```

35      extends ←----- 3'      5'
                        GGGGGGGGGG Primer 1
      ....ZZZZZZZZZZZZZZZZAAAAAAAAAAXXXXXXXXXXCCCCCCCCCZZZZZZZZZZZZZZZZ....
40      original template strand+

      original template strand-
      ....ZZZZZZZZZZZZZZZZTTTTTTTTTTYYYYYYYYYYGGGGGGGGGGZZZZZZZZZZZZZZZZ....
45      Primer 2      AAAAAA AAAA -----> extends
                        5'      3'

```

[0056] On denaturation of the two duplexes formed, the products are:

3' 5'
zzzzzzzzzzzzzzzzTTTTTTTTTTTTYYYYYYYYYYGGGGGGGGGG
 newly synthesized long product 1

5' 3'
zzzzzzzzzzzzzzzzAAAAAAAAXXXXXXXXXXXCCCCCCCCCzzzzzzzzzzzzzzzz....
 original template strand⁺

3' 5'
zzzzzzzzzzzzzzzzTTTTTTTTTTTTYYYYYYYYYYGGGGGGGGGGzzzzzzzzzzzzzzzz....
 original template strand⁻

5' 3'
 AAAAAAAAAXXXXXXXXXXXCCCCCCCCCzzzzzzzzzzzzzzzz....
 newly synthesized long product 2

[0057] If these four strands are allowed to rehybridize with Primers 1 and 2 in the next cycle, agent for polymerization will catalyze the following reactions:

Primer 2 5' AAAAAAAA → extends to here

3'....zzzzzzzzzzzzzzzzTTTTTTTTTTTTYYYYYYYYYYGGGGGGGGGG 5'
 newly synthesized long product 1

extends ← GGGGGGGGGG 5' Primer 1

5'....zzzzzzzzzzzzzzzzAAAAAAAAXXXXXXXXXXXCCCCCCCCCzzzzzzzzzzzzzzzz....3'
 original template strand⁺

Primer 2 5' AAAAAAAA → extends

3'....zzzzzzzzzzzzzzzzTTTTTTTTTTTTYYYYYYYYYYGGGGGGGGGGzzzzzzzzzz....5'
 original template strand⁻

extends to here ← GGGGGGGGGG 5' Primer 1

5' AAAAAAAAAXXXXXXXXXXXCCCCCCCCCzzzzzzzzzzzzzzzz....3'
 newly synthesized long product 2

[0058] If the strands of the above four duplexes are separated, the following strands are found:

5' AAAAAAAAAAXXXXXXXXXXCCCCCCCCC 3'
newly synthesized [S⁺]

3'....zzzzzzzzzzzzzzzzzzTTTTTTTTTYYYYYYYYYGGGGGGGGG 5'
first cycle synthesized long product 1

3'....zzzzzzzzzzzzzzzzzzTTTTTTTTTYYYYYYYYYGGGGGGGGG 5'
newly synthesized long product 1

5'....zzzzzzzzzzzzzzzzzzAAAAAAAAAXXXXXXXXXXCCCCCCCCCzzzzzzzzzz....3'
original template strand⁺

5' AAAAAAAAAAXXXXXXXXXXCCCCCCCCCzzzzzzzzzzzzzzzz....3'
newly synthesized long product 2

3'..zzzzzzzzzzzzzzzzTTTTTTTTTYYYYYYYYYGGGGGGGGGzzzzzzzzzzzzzzzz....5'
original template strand⁻

3' TTTTTTTTTTYYYYYYYYYGGGGGGGGG 5'
newly synthesized [S⁻]

5' AAAAAAAAAAXXXXXXXXXXCCCCCCCCCzzzzzzzzzzzzzzzz....3'
first cycle synthesized long product 2

[0059] It is seen that each strand which terminates with the oligonucleotide sequence of one primer and the complementary sequence of the other is the specific nucleic acid sequence [S] that is desired to be produced.

[0060] The steps of this process can be repeated indefinitely, being limited only by the amount of Primers 1 and 2, the agent for polymerization and nucleotides present. For detection, the number of cycles used is that required to produce a detectable signal, an amount which will depend, e.g., on the nature of the sample. For example, if the sample is pure or diluted, fewer cycles may be required than if it is a complex mixture. If the sample is human genomic DNA, preferably the number of cycles is from about 10-30.

[0061] The amount of original nucleic acid remains constant in the entire process, because it is not replicated. The amount of the long products increases linearly because they are produced only from the original nucleic acid. The amount of the specific sequence increases exponentially. Thus, the specific sequence will become the predominant species. This is illustrated in the following table, which indicates the relative amounts of the species theoretically present after n cycles, assuming 100% efficiency at each cycle:

Number of Double Strands After 0 to n Cycles			
Cycle Number	Template	Long Products	Specific Sequence [S]
0	1	-	-
1	1	1	0
2	1	2	1
3	1	3	4
5	1	5	26
10	1	10	1013
15	1	15	32,752
20	1	20	1,048,555
n	1	n	(2 ⁿ -n-1)

When a single-stranded nucleic acid is utilized as the template, only one long product is formed per cycle.

[0062] The kit herein may be utilized to clone a particular nucleic acid sequence for insertion into a suitable expression vector. The vector may then be used to transform an appropriate host organism to produce the gene product of the sequence by standard methods of recombinant DNA technology.

[0063] Normally, such cloning would either involve direct ligation into a vector or the addition of oligonucleotide linkers

followed by restriction enzyme cleavage. Both of these methods involve, however, the inefficient blunt-end ligation reaction. Also, neither technique would control for the orientation or multiplicity of insertion of the amplified product into the cloning vector.

[0064] The amplification process carried out by using the kit herein may yield a mixture of nucleic acids, resulting from the original template nucleic acid, the expected target amplified products, and various background non-target products. The amplified product can also be a mixture if the original template DNA contains multiple target sequences, such as in a heterozygous diploid genome or when there is a family of related genes.

[0065] The primers herein may be modified to assist the rapid and specific cloning of the mixture of DNAs produced by the amplification reaction. In such modification the same or different restriction sites are incorporated at the 5' ends of the primers to result in restriction sites at the two ends of the amplified product. When cut with the appropriate enzymes, the amplified product can then be easily inserted into plasmid or viral vectors and cloned. This cloning allows the analysis or expression of individual amplified products, not a mixture.

[0066] Although the same restriction site can be used for both primers, the use of different sites allows the insertion of the product into the vector with a specific orientation and suppresses multiple insertions as well as insertions arising from amplifications based on only one of the two primers. The specific orientation is useful when cloning into single-strand sequencing vectors, when single-strand hybridization probes are used, or when the cloned product is being expressed.

[0067] One method to prepare the primers is to choose a primer sequence which differs minimally from the target sequence. Regions in which each of the primers is to be located are screened for homology to restriction sites appropriate to the desired vector. For example, the target sequence "CAGTATCCGA..." differs by only one base from one containing a *Bam*HI site. A primer sequence is chosen to match the target exactly at its 3' end, and to contain the altered sequence and restriction site near its 5' end (for example, "CAGgATCCGA...", where the lower case letter symbolizes a mismatch with the target sequence). This minimally altered sequence will not interfere with the ability of the primer to hybridize to the original target sequence and to initiate polymerization. After the first amplification cycle the primer is copied, becomes the target, and matches exactly with new primers. After the amplification process, the products are cleaved with the appropriate restriction enzymes, optionally separated from inhibitors of ligation such as the nucleotide triphosphates and salts by passing over a desalting column or molecular weight chromatography column, and inserted by ligation into a cloning vector such as bacteriophage M13. The gene may then be sequenced and/or expressed using well known techniques.

[0068] The second method for preparing the primers involves taking the 3' end of the primers from the target sequence and adding the desired restriction site(s) to the 5' end of the primer. For the above example, a *Hind*III site could be added to make the sequence "cgaagcttCAGTATCCGA...", where lower case letters are as described above. The added bases would not contribute to the hybridization in the first cycle of amplification, but would match in subsequent cycles. The final amplified products are then cut with restriction enzyme(s) and cloned and expressed as described above. The gene being amplified may be, for example, human beta-hemoglobin or the human HLA DQ, DR or DP- α and - β genes.

[0069] In addition, the kit herein can be used for *in vitro* mutagenesis. The oligodeoxyribonucleotide primers need not be exactly complementary to the DNA sequence which is being amplified. It is only necessary that they be able to hybridize to the sequence sufficiently well to be extended by the polymerase enzyme or by whatever other inducing agent is employed. The product of a polymerase chain reaction wherein the primers employed are not exactly complementary to the original template will contain the sequence of the primer rather than the template, thereby introducing an *in vitro* mutation. In further cycles this mutation will be amplified with an undiminished efficiency because no further mispaired primings are required. The mutant thus produced may be inserted into an appropriate vector by standard molecular biological techniques and might confer mutant properties on this vector such as the potential for production of an altered protein.

[0070] The process of making an altered DNA sequence as described above could be repeated on the altered DNA using different primers so as to induce further sequence changes. In this way a series of mutated sequences could gradually be produced wherein each new addition to the series could differ from the last in a minor way, but from the original DNA source sequence in an increasingly major way. In this manner changes could be made ultimately which were not feasible in a single step due to the inability of a very seriously mismatched primer to function.

[0071] In addition, the primer can contain as part of its sequence a non-complementary sequence provided that a sufficient amount of the primer contains a sequence which is complementary to the strand to be amplified. For example, a nucleotide sequence which is not complementary to the template sequence (such as, e.g., a promoter, linker, coding sequence, etc.) may be attached at the 5' end of one or both of the primers, and thereby appended to the product of the amplification process. After the extension primer is added, sufficient cycles are run to achieve the desired amount of new template containing the non-complementary nucleotide insert. This allows production of large quantities of the combined fragments in a relatively short period of time (e.g., two hours or less) using a simple technique.

[0072] Moreover, the kit herein may be used to synthesize a nucleic acid fragment from an existing nucleic acid

fragment which is shorter than its product (called the core segment) using certain primers the 3' ends of which are complementary to or substantially complementary to the 3' ends of the single strands produced by separating the strands of the original shorter nucleic acid fragments, and the 5' ends of which primers contain sequence information to be appended to the core segment. This process comprises:

(a) treating the strands of said existing fragment with two oligonucleotide primers under conditions such that an extension product of each primer is synthesized which is complementary to each nucleic acid strand, wherein said primers are selected so as to be substantially complementary to the 3' end of each strand of said existing fragment such that the extension product synthesized from one primer, when it is separated from its complement, can serve as a template for synthesis of the extension product of the other primer, and wherein each primer contains, at its 5' end, a sequence of nucleotides which are not complementary to said existing fragment and which correspond to the two ends of the nucleic acid fragment being synthesized;

(b) separating the primer extension products from the templates on which they were synthesized to produce single-stranded molecules; and

(c) treating the single-stranded molecules generated from step (b) with the primers of step (a) under conditions such that a primer extension product is synthesized using each of the single strands produced in step (b) as a template so as to produce two intermediate double-stranded nucleic acid molecules, into each of which has been incorporated the nucleotide sequence present in the 5' end of one of the oligonucleotide primers, and two full-length double-stranded nucleic acid molecules, into each of which has been incorporated the nucleotide sequence present in the 5' ends of both of the oligonucleotide primers;

(d) repeating steps (b) and (c) for a sufficient number of times to produce the full-length double-stranded molecules in an effective amount;

(e) treating the strands of the product of step (d) with two primers so as to lengthen the product of step (d) on both ends; and

(f) repeating steps (a)-(d) using the product of step (d) as the core fragment and two oligonucleotide primers which are complementary or substantially complementary to the 3' ends of the single strands produced by separating the strands of the product of step (d).

[0073] Steps (b) and (c) are repeated as often as necessary, usually at least 5 times, to produce the required amount of the full-length double-stranded product to synthesize the final product (i.e., the effective amount). In addition, the core segment may be obtained as the product of a previous amplification cycle. The product produced in step (d) may be purified before a new cycle of extension and amplification, or used directly by employing the reaction mixture containing the product.

[0074] If the 3' ends of the primers are not exactly complementary to the 3' ends of the single strands of the original shorter nucleic acid, the core fragment of the product will not be exactly the same as the sequence information resident in the original shorter nucleic acid. Therefore, mutants of the original nucleic acid may be made by using primers which are substantially complementary at their 3' ends to the 3' ends of the single strands of the original shorter nucleic acid.

[0075] If restriction site linkers are incorporated into the primers, then the amplified double-stranded products can be digested with the appropriate restriction enzymes and ligated directly into an M13 vector for rapid cloning and sequencing. The M13 plaques containing the specific amplified target sequences can be identified by hybridizing plaque lift filters with a probe specific for the target sequence.

[0076] The kit herein may also be used to enable detection and/or characterization of specific nucleic acid sequences associated with infectious diseases, genetic disorders or cellular disorders such as cancer, e.g., oncogenes. Amplification is useful when the amount of nucleic acid available for analysis is very small, as, for example, in the prenatal diagnosis of sickle cell anemia using DNA obtained from fetal cells. Amplification is particularly useful if such an analysis is to be done on a small sample using non-radioactive detection techniques which may be inherently insensitive, or where radioactive techniques are being employed but where rapid detection is desirable.

[0077] For purposes of this invention genetic diseases may include specific deletions and/or mutations in genomic DNA from any organism, such as, e.g., sickle cell anemia, cystic fibrosis, α -thalassemia, β -thalassemia, and the like. Sickle cell anemia can be readily detected via oligomer restriction analysis or a RFLP-like analysis following amplification of the appropriate DNA sequence by the present method. α -Thalassemia can be detected by the absence of a sequence, and β -thalassemia can be detected by the presence of a polymorphic restriction site closely linked to a mutation which causes the disease.

[0078] All of these genetic diseases may be detected by amplifying the appropriate sequence and analyzing it by Southern blots without using radioactive probes. In such a process, for example, a small sample of DNA from, e.g., amniotic fluid containing a very low level of the desired sequence is amplified, cut with a restriction enzyme, and analyzed via a Southern blotting technique. The use of non-radioactive probes is facilitated by the high level of the amplified signal.

[0079] In another embodiment a small sample of DNA may be amplified to a convenient level and then a further cycle of extension reactions performed wherein nucleotide derivatives which are readily detectable (such as ^{32}P -labeled or biotin labeled nucleoside triphosphates) are incorporated directly into the final DNA product, which may be analyzed by restriction and electrophoretic separation or any other appropriate method. An example of this technique in a model system is demonstrated in Figure 5.


[0080] In a further embodiment, demonstrated in a model system in Figure 3, the nucleic acid may be exposed to a particular restriction endonuclease prior to amplification. Since a sequence which has been cut cannot be amplified, the appearance of an amplified fragment, despite prior restriction of the DNA sample, implies the absence of a site for the endonuclease within the amplified sequence. The presence or absence of an amplified sequence can be detected by an appropriate method.

[0081] A practical application of this technique can be illustrated by its use in facilitating the detection of sickle cell anemia via the oligomer restriction technique described herein and in Saiki et al., *Biotechnology*, 3:1008-1012 (1985). Sickle cell anemia is a hemoglobin disease which is caused by a single base pair change in the sixth codon of the β -globin gene. Figure 6 illustrates the sequences of normal and sickle cell β -globin genes in the region of their polymorphism, where the single bars mark the location of a *DdeI* site present only in the normal gene and where the double bars mark the location of a *HinfI* site which is non-polymorphic and thus present in both the normal and sickle cell alleles. Figure 7 illustrates the process of oligomer restriction of normal β -globin DNA using a probe spanning both restriction sites and labeled where the asterisk appears. (The probe is preferably labeled at the end which is fewer base pairs from the restriction site than the other end of the probe.) The DNA, amplified as provided herein, is denatured and annealed to the labeled probe. The amplification may be carried out at elevated temperatures (35-40°C) in the presence of dimethyl sulfoxide to minimize formation of secondary structure. The enzyme *DdeI* cleaves the DNA at the reformed *DdeI* site and generates a labeled octamer. Under the conditions used in the test the octamer is short enough to dissociate from the duplex. The subsequent addition of the enzyme *HinfI* has no effect on the now single-stranded octamer. Figure 8 illustrates the same process applied to the sickle cell allele of β -globin DNA. The enzyme *DdeI* cannot cleave the duplex formed by the amplified DNA and the labeled probe because of the A-A base pair mismatch. The enzyme *HinfI*, however, does restrict the hybrid and a labeled trimer is produced. In practice the method can diagnose the DNA of an individual as being either homozygous for the wild type, homozygous for the sickle type or a heterozygous carrier of the sickle cell trait, since a specific signal is associated with the presence of either allele. Use of this above-described method to amplify the pertinent sequence allows for a rapid analysis of a single copy gene using a probe with only a single ^{32}P label.


[0082] Various infectious diseases can be diagnosed by the presence in clinical samples of specific DNA sequences characteristic of the causative microorganism. These include bacteria, such as *Salmonella*, *Chlamydia*, *Neisseria*; viruses, such as the hepatitis viruses, and parasites, such as the *Plasmodium* responsible for malaria. U.S. Patent 4,358,535 issued to Falkow describes the use of specific DNA hybridization probes for the diagnosis of infectious diseases. A problem inherent in the Falkow procedure is that a relatively small number of pathogenic organisms may be present in a clinical sample from an infected patient and the DNA extracted from these may constitute only a very small fraction of the total DNA in the sample. Specific amplification of suspected sequences prior to immobilization and hybridization detection of the DNA samples could greatly improve the sensitivity and specificity of these procedures.

[0083] Routine clinical use of DNA probes for the diagnosis of infectious diseases would be simplified considerably if non-radioactively labeled probes could be employed as described in EP 63,879 to Ward. In this procedure biotin-containing DNA probes are detected by chromogenic enzymes linked to avidin or biotin-specific antibodies. This type of detection is convenient, but relatively insensitive. The combination of specific DNA amplification by the present method and the use of stably labeled probes could provide the convenience and sensitivity required to make the Falkow and Ward procedures useful in a routine clinical setting.

[0084] In addition, the probe may be a biotinylated probe in which the biotin is attached to a spacer arm of the formula:



where Y is O, NH or N-CHO, x is a number from 1 to 4, and y is a number from 2 to 4. The spacer arm is in turn attached to a psoralen moiety of the formula:



The psoralen moiety intercalates into and crosslinks a "gapped circle probe as described by Courage-Tebbe et al., *Biochim. Biophys. Acta*, 697 (1982) 1-5, wherein the single-stranded hybridization region of the gapped circle spans the region contained in the primers.

[0085] The kit can also be utilized to produce sufficient quantities of DNA from a single copy human gene such that detection by a simple non-specific DNA stain such as ethidium bromide can be employed so as to make a DNA diagnosis directly.

[0086] In addition to detecting infectious diseases and pathological abnormalities in the genome of organisms, the kit herein can also be used to detect DNA polymorphism which may not be associated with any pathological state.

[0087] The following examples are offered by way of illustration and are not intended to limit the invention in any manner. In these examples all percentages are by weight if for solids and by volume if for liquids, and all temperatures are in degrees Celsius unless otherwise noted.

EXAMPLE 1

[0088] A 25 base pair sequence having the nucleotide sequence

5' CCTCGGCACCGTCACCGTGGATGCT 3'
3' GGAGCCGTGGCAGTGGGACCTACGA 5'

contained on a 47 base pair *FokI* restriction fragment of pBR322 obtainable from ATCC was prepared as follows. A *FokI* digest of pBR322 containing the 47-bp fragment was produced by digesting pBR322 with *FokI* in accordance with the conditions suggested by the supplier, New England Biolabs Inc. The primers which were utilized were 5' d(CCTCG-GCACCG) 3' and 5' d(AGCATCCAGGGTG) 3', and were prepared using conventional techniques. The following ingredients were added to 33 μ l of buffer which consisted of 25 mM potassium phosphate, 10 mM magnesium chloride and 100 mM sodium chloride at pH 7.5: 2433 pmoles of each of the primers described above, 2.4 pmoles of the *FokI* digest of pBR322, 12 nmoles of dATP, 22 nmoles of dCTP, 19 nmoles of dGTP and 10 nmoles of TTP.

[0089] The mixture was heated to 85°C for five minutes and allowed to cool to ambient temperature. Five units of the Klenow fragment of *E. coli* DNA polymerase I were added and the temperature was maintained for 15 minutes. After that time, the mixture was again heated to 85°C for five minutes and allowed to cool. Five units of the Klenow fragment were again added and the reaction was carried out for 15 minutes. The heating, cooling and synthesis steps were repeated eleven more times.

[0090] After the final repetition, a 5 μ l aliquot was removed from the reaction mixture. This was heated to 85°C for three minutes and allowed to cool to ambient temperature. 12.5 pmoles of α -³²P-deoxycytidine triphosphate and 5 units of Klenow fragment were added and the reaction was allowed to proceed for 15 minutes. The labeled products were examined by polyacrylamide gel electrophoresis. The *FokI* digest was labeled in a similar fashion and served as a control and molecular weight markers. The only heavily labeled band visible after the 13 cycles was the intended 25 base pair sequence.

EXAMPLE 2

[0091] The desired sequence to be amplified was a 94 base pair sequence contained within the human beta-globin gene and spanning the *MstII* site involved in sickle cell anemia. The sequence has the nucleotide sequence shown in Figure 1.

I. Synthesis of Primers

[0092] The following two oligodeoxyribonucleotide primers were prepared by the method described below:

5' CACAGGGCAGTAACG 3' Primer A

and

5' TTTGCTTCTGACACA 3' Primer B

[0093] Automated Synthesis Procedures: The diethylphosphoramidites, synthesized according to Beaucage and Caruthers (Tetrahedron Letters (1981) 22:1859-1862) were sequentially condensed to a nucleoside derivatized controlled pore glass support using a Biosearch SAM-1. The procedure included detritylation with trichloroacetic acid in dichloromethane, condensation using benzotriazole as activating proton donor, and capping with acetic anhydride and dimethylaminopyridine in tetrahydrofuran and pyridine. Cycle time was approximately 30 minutes. Yields at each step were essentially quantitative and were determined by collection and spectroscopic examination of the dimethoxytrityl alcohol released during detritylation.

[0094] Oligodeoxyribonucleotide Deprotection and Purification Procedures: The solid support was removed from the column and exposed to 1 ml concentrated ammonium hydroxide at room temperature for four hours in a closed tube. The support was then removed by filtration and the solution containing the partially protected oligodeoxynucleotide was brought to 55°C for five hours. Ammonia was removed and the residue was applied to a preparative polyacrylamide gel. Electrophoresis was carried out at 30 volts/cm for 90 minutes after which the band containing the product was identified by UV shadowing of a fluorescent plate. The band was excised and eluted with 1 ml distilled water overnight at 4°C. This solution was applied to an Altech RP18 column and eluted with a 7-13% gradient of acetonitrile in 1% ammonium acetate buffer at pH 6.0. The elution was monitored by UV absorbance at 260 nm and the appropriate fraction collected, quantitated by UV absorbance in a fixed volume and evaporated to dryness at room temperature in a vacuum centrifuge.

[0095] Characterization of Oligodeoxyribonucleotides: Test aliquots of the purified oligonucleotides were ³²P labeled with polynucleotide kinase and γ -³²P-ATP. The labeled compounds were examined by autoradiography of 14-20% polyacrylamide gels after electrophoresis for 45 minutes at 50 volts/cm. This procedure verifies the molecular weight. Base composition was determined by digestion of the oligodeoxyribonucleotide to nucleosides by use of venom diesterase and bacterial alkaline phosphatase and subsequent separation and quantitation of the derived nucleosides using a reverse phase HPLC column and a 10% acetonitrile, 1% ammonium acetate mobile phase.

II. Source of DNA

A. Extraction of Whole Human Wild-Type DNA

[0096] Human genomic DNA homozygous for normal β -globin was extracted from the cell line Molt4 (obtained from Human Genetic Mutant Cell Repository and identified as GM2219c) using the technique described by Stetler et al., Proc. Nat. Acad. Sci. (1982), 79:5956-5970.

B. Construction of Cloned Globin Genes

[0097] A 1.9 kb BamHI fragment of the normal β -globin gene was isolated from the cosmid pFC11 and inserted into the BamHI site of pBR328 (Soberon, et al., Gene (1980) 9:287-305). This fragment, which encompasses the region that hybridizes to the synthetic 40-mer probe, includes the first and second exons, first intron, and 5' flanking sequences of the gene (Lawn et al., Cell (1978), 15:1157-1174). This clone was designated pBR328:HbA and deposited under ATCC No. 39,698 on May 25, 1984.

[0098] The corresponding 1.9 kb BamHI fragment of the sickle cell allele of β -globin was isolated from the cosmid pFC12 and cloned as described above. This clone was designated pBR328:HbS and deposited under ATCC No. 39,699 on May 25, 1984.

[0099] Each recombinant plasmid was transformed into and propagated in E. coli MM294 (ATCC No. 39,607).

C. Digestion of Cloned Globin Genes with MstII

[0100] A total of 100 μ g each of pBR328:HbA and pBR328:HbS were individually digested with 20 units of MstII (New England Biolabs) for 16 hours at 37°C in 200 μ l of 150 mM NaCl, 12 mM Tris HCl (pH 7.5), 12 mM MgCl₂, 1 mM dithiothreitol (DTT), and 100 μ g/ml bovine serum albumin (BSA). The products are designated pBR328:HbA/MstII and pBR328:HbS/MstII, respectively.

III. Polymerase Chain Reaction

[0101] To 100 µl of buffer consisting of 60 mM sodium acetate, 30 mM Tris acetate and 10 mM magnesium acetate at pH 8.0 was added 2 µl of a solution containing 100 picomoles of Primer A (of the sequence d(CACAG-GGCACTAACG)), 100 picomoles of Primer B (of the sequence d(TTTGCTTCTGACACA)) and 1000 picomoles each of dATP, dCTP, dGTP and TTP. In addition, one of the following sources of DNA described above was added:

10 µg whole human wild-type DNA (Reaction I)

0.1 picomole pBR328:HbA (Reaction II)

0.1 picomole pBR328:HbS (Reaction III)

0.1 picomole pBR328:HbA/MstII (Reaction IV)

0.1 picomole pBR328:HbS/MstII (Reaction V)

No target DNA (Reaction VI)

[0102] Each resulting solution was heated to 100°C for four minutes and allowed to cool to room temperature for two minutes, whereupon 1 µl containing four units of Klenow fragment of *E. coli* DNA polymerase was added. Each reaction was allowed to proceed for 10 minutes, after which the cycle of adding the primers, nucleotides and DNA, heating, cooling, adding polymerase, and reacting was repeated nineteen times for Reaction I and four times for Reactions II-VI.

[0103] Four microliter aliquots of Reactions I and II removed before the first cycle and after the last cycle of each reaction were applied to a 12% polyacrylamide gel 0.089 M in Tris-borate buffer at pH 8.3 and 2.5 mM in EDTA. The gel was electrophoresed at 25 volts/cm for four hours, transferred to a nylon membrane serving as solid phase support and probed with a 5'-³²P-labeled 40 bp synthetic fragment, prepared by standard techniques, of the sequence

5' d (TCCTGAGGAGAAGTCTGCCGTTACTGCCCTGTGGGGCAAG) 3'

in 30% formamide, 3 x SSPE, 5 x Denhardt's, 5% sodium dodecyl sulfate at pH 7.4. Figure 2 is an autoradiograph of the probed nylon membrane for Reactions I and II. Lane 1 is 0.1 picomole of a 58-bp synthetic fragment control one strand of which is complementary to the above probe. Lane 2 is 4 µl of Reaction I prior to the first amplification cycle. Lane 3 is 4 µl of Reaction I after the 20th amplification cycle. Lane 4 is 4 µl of Reaction II after five amplification cycles. Lane 5 is a molecular weight standard consisting of a *FokI* (New England Biolabs) digest of pBR322 (New England Biolabs) labeled with alpha-³²P-dNTPs and polymerase. Lane 3 shows that after twenty cycles the reaction mixture I contained a large amount of the specific sequence of the proper molecular weight and no other detectable products. Reaction mixture II after five cycles also contained this product, as well as the starting nucleic acid and other products, as shown by Lane 4.

[0104] To 5.0 µl aliquots of Reactions II-VI after the fourth cycle were added 5 pmoles of each primer described above. The solutions were heated to 100°C for four minutes and allowed to equilibrate to room temperature. Three pmoles each of alpha-³²P-dATP, alpha-³²P-dCTP, alpha-³²P-dGTP and alpha-³²P-TTP and four units of Klenow fragment were added. The reaction, in a final volume of 10 µl and at the salt concentrations given above, was allowed to proceed for 10 minutes. The polymerase activity was terminated by heating for 20 minutes at 60°C. Four µl aliquots of Reactions II-VI were loaded onto a 12% polyacrylamide gel 0.089 M in Tris-borate buffer at pH-8.3 and 2.5 mM in EDTA. The gel was electrophoresed at 25 volts/cm for four hours after which autoradiography was performed.

[0105] Figure 3 is an autoradiograph of the electrophoresis. Lane 1 is a molecular weight standard, Lane 2 is Reaction II, Lane 3 is Reaction III, Lane 4 is Reaction IV and Lane 5 is Reaction V. Another lane for Reaction VI with no DNA as control had no images in any of the lanes. It can be seen from the figure that the 94-bp fragment predicted from the target DNA was present only where intact β-globin DNA sequences were available for amplification, i.e., pBR328:HbA (Lane 2), pBR328:HbS (Lane 3) and pBR328:HbS/MstII (Lane 5). *MstII* digestion cuts pBR328:HbA in the 94-mer sequence rendering it incapable of being amplified, and the 94-mer band does not appear in Lane 4. In contrast, the 94-mer sequence in pBR328:HbS does not cut when the plasmid is digested with *MstII* and thus is available for amplification as shown in Lane 5.

[0106] Figure 4 illustrates the chain reaction for three cycles in amplifying the 94-bp sequence. PC01 and PC02 are Primers A and B. The numbers on the right indicate the cycles, whereas the numbers on the left indicates the cycle

number in which a particular molecule was produced.

EXAMPLE 3

5 **[0107]** This example illustrates amplification of a 110 bp sequence spanning the allelic MstII site in the human hemoglobin gene.

[0108] A total of 1.0 microgram whole human DNA, 100 picomoles d(ACACAAGTGTGTTCACTAGC) and 100 picomoles d(CAACTTCATCCACGTTCAACC) the primers having been prepared by the technique of Example 2, were dissolved in 100 µl of a solution which was:

10

1.5 mM in each of the four deoxyribonucleoside triphosphates

30 mM in Tris acetate buffer at pH 7.9

15

60 mM in sodium acetate

10 mM in magnesium acetate

0.25 mM in dithiothreitol

20

[0109] The solution was heated to 100°C for one minute and brought rapidly to 25°C for one minute, after which was added 2.5 units Klenow fragment of DNA polymerase. The polymerase reaction was allowed to proceed for two minutes at 25°C, after which the cycle of heating, cooling, adding Klenow, and reacting was repeated as often as desired.

25

[0110] With a 70% efficiency at each cycle, 15 cycles resulted in the synthesis of 1.4 femtomoles of the desired 110 bp fragment of the β-globin gene.

EXAMPLE 4

30 **[0111]** This example illustrates amplification of a 240 bp sequence spanning the allelic MstII site in the human hemoglobin gene. This sequence contains NcoI, HinfI and MstII restriction sites.

[0112] To 100 µl of a mixture of 60 mM sodium acetate, 30 mM Tris acetate and 10 mM magnesium acetate at pH 8.0 containing 0.1 pmole pBR328:HbA was added 2 µl of Solution A containing:

35

100 pmoles d(GGTGGCCAATCTACTCCAGG) primer

100 pmoles d(TAACCTTGATACCAACCTGCCC) primer

40

1000 pmoles each of dATP, dCTP, dGTP and TTP

45

[0113] The two primers were prepared by the technique described in Example 2. The solution was heated to 100°C for four minutes and allowed to cool in ambient air for two minutes, after which was added 1 µl containing four units Klenow fragment of E. coli DNA polymerase. The reaction was allowed to proceed for 10 minutes after which the cycle of solution A addition, heating, cooling, adding polymerase, and reacting was repeated three times. To a 5.0 µl aliquot of the reactions was added 5 picomoles of each oligonucleotide primer described above. The solution was heated to 100°C for four minutes and allowed to come to ambient temperature, after which 3 picomoles each of the alpha-³²P-labeled deoxyribonucleoside triphosphates and 4 units Klenow fragment were added. The reaction, in a final volume of 10 µl and at the salt concentrations given above, was allowed to proceed for 10 minutes. The polymerase activity was terminated by heating for 20 minutes at 60°C. Two µl aliquots were digested with NcoI, MstII, or HinfI and loaded onto a 12% polyacrylamide gel 0.089 M in Tris-borate buffer at pH 8.3 and 2.5 mM in EDTA. The gel was electrophoresed at 25 volts/cm for four hours and autoradiography was performed. Figure 5 illustrates the autoradiograph of the electrophoresis, where Lane 1 is the molecular weight standard, Lane 2 is without digestion with enzyme (240 bp intact), Lane 3 is digestion with NcoI (131 and 109 bp), Lane 4 is digestion with MstII (149 and 91 bp), and Lane 5 is digestion with HinfI (144 and 96 bp). The autoradiograph is consistent with the amplification of the 240 bp sequence.

55

EXAMPLE 5

[0114] This example illustrates use of the process herein to detect sickle cell anemia by sequential digestion.

5 Synthesis and Phosphorylation of Oligodeoxyribonucleotides

[0115] A labeled DNA probe, RS06, of the sequence:

10 5' *CTGACTCCTGAGGAGAAGTCTGCCGTTACTGCCCTGTGGG 3'

where * indicates the label, and an unlabeled blocking oligomer, RS10, of the sequence

15 3' GACAGAGGTCACCTCTTCAGACGGCAATGACGGGACACCC 5'

which has three base pair mismatches with RS06 were synthesized according to the procedures provided in Example 2(I) The probe RS06 was labeled by contacting five pmole thereof with 4 units of T4 polynucleotide kinase (New England Biolabs) and 50 pmole γ -³²P-ATP (New England Nuclear, about 7200 Ci/mmole) in a 40 μ l reaction volume containing 70 mM Tris buffer (pH 7.6), 10 mM MgCl₂, 1.5 mM spermine, and 2.5 mM dithiothreitol for 90 minutes at 37°C. The total volume was then adjusted to 100 μ l with 25 mM EDTA and purified according to the procedure of Maniatis et al., Molecular Cloning (1982), 464-465 over a 1 ml Bio Gel P-4 spin dialysis column from BioRad equilibrated with Tris-EDTA (TE) buffer (10 mM Tris buffer, 0.1 mM EDTA, pH 8.0). The labeled probe was further purified by electrophoresis on a 18% polyacrylamide gel (19:1 acrylamide:BIS, BioRad) in Tris-boric acid-EDTA (TBE) buffer (89 mM Tris, 89 mM boric acid, 2.5 mM EDTA, pH 8.3) for 500 vhr. After localization by autoradiography, the portion of the gel containing the labeled probe was excised, crushed and eluted into 0.2 ml TE buffer overnight at 4°C. TCA precipitation of the reaction product indicated that the specific activity was 4.9 Ci/mmole and the final concentration was 20 pmole/ml.

[0116] The unlabeled RS10 blocking oligomer was used at a concentration of 200 pmole/ml.

30 Isolation of Human Genomic DNA from Cell Lines

[0117] High molecular weight genomic DNA was isolated from the lymphoid cell lines Molt4, SC-1 and GM2064 using essentially the method of Stetler et al., PNAS (1982), 79, 5966-5979 (for Molt4) and Maniatis et al., Molecular Cloning (1982), 280-281.

[0118] Molt4 (Human Mutant Cell Repository, GM2219C) is a T cell line homozygous for normal β -globin, and SC-1, deposited with ATCC on March 19, 1985, is an EBV-transformed B cell line homozygous for the sickle cell allele. GM2064 (Human Mutant Cell Repository, GM2064) was originally isolated from an individual homozygous for hereditary persistence of fetal hemoglobin (HPFH) and contains no beta- or delta-globin gene sequences. All cell lines were maintained in RPMI-1640 with 10% fetal calf serum.

40 Isolation of Human Genomic DNA from Clinical Blood Samples

[0119] A clinical blood sample designated CH12 from a known sickle cell carrier (AS) was obtained from Dr. Bertram Lubin of Children's Hospital in Oakland, California. Genomic DNA was prepared from the buffy coat fraction, which is composed primarily of peripheral blood lymphocytes, using a modification of the procedure described by Nunberg et al., Proc. Nat. Acad. Sci., 75, 5553-5556 (1978).

[0120] The cells were resuspended in 5 ml Tris-EDTA-NaCl (TEN) buffer (10 mM Tris buffer pH 8, 1 mM EDTA, 10 mM NaCl) and adjusted to 0.2 mg/ml proteinase K, 0.5% SDS, and incubated overnight at 37°C. Sodium perchlorate was then added to 0.7 M and the lysate gently shaken for 1-2 hours at room temperature. The lysate was extracted with 30 ml phenol/chloroform (1:1), then with 30 ml chloroform, and followed by ethanol precipitation of the nucleic acids. The pellet was resuspended in 2 ml of TE buffer and RNase A added to 0.005 mg/ml. After digestion for one hour at 37°C, the DNA was extracted once each with equal volumes of phenol, phenol/chloroform, and chloroform, and ethanol precipitated. The DNA was resuspended in 0.5 ml TE buffer and the concentration was determined by absorbance at 260 nm.

Polymerase Chain Reaction to Amplify Selectively β -Globin Sequences

[0121] Two micrograms of genomic DNA was amplified in an initial 100 μ l reaction volume containing 10 mM Tris buffer (pH 7.5), 50 mM NaCl, 10 mM $MgCl_2$, 150 pmole of Primer A of the sequence d(CACAGGGCACTAACG), and 150 pmole of Primer B of the sequence d(CTTGCTTCTGACACA) and overlaid with about 100 μ l mineral oil to prevent evaporation.

[0122] Each DNA sample underwent 15 cycles of amplification where one cycle is composed of three steps:

1) Denature in a heat block set at 95°C for two minutes.

2) Transfer immediately to a heat block set at 30°C for two minutes to allow primers and genomic DNA to anneal.

3) Add 2 μ l of a solution containing 5 units of the Klenow fragment of *E. coli* DNA polymerase I (New England Biolabs), 1 nmole each of dATP, dCTP, dGTP and TTP, in a buffer composed of 10 mM Tris (pH 7.5), 50 mM NaCl, 10 mM $MgCl_2$, and 4 mM dithiothreitol. This extension reaction was allowed to proceed for 10 minutes at 30°C.

[0123] After the final cycle, the reaction was terminated by heating at 95°C for two minutes. The mineral oil was extracted with 0.2 ml of chloroform and discarded. The final reaction volume was 130 μ l.

Hybridization/Digestion of Amplified Genomic DNA with Probes and DdeI/HinfI

[0124] Forty-five microliters of the amplified genomic DNA was ethanol precipitated and resuspended in an equal volume of TE buffer. Ten microliters (containing the pre-amplification equivalent of 154 ng of genomic DNA) was dispensed into a 1.5 ml Microfuge tube and 20 μ l of TE buffer to a final volume of 30 μ l. The sample was overlaid with mineral oil and denatured at 95°C for 10 minutes. Ten microliters of 0.6 M NaCl containing 0.02 pmole of labeled RS06 probe was added to the tube, mixed gently, and immediately transferred to a 56°C heat block for one hour. Four microliters of unlabeled RS10 blocking oligomer (0.8 pmole) was added and the hybridization continued for an additional 10 minutes at the same temperature. Five microliters of 60 mM $MgCl_2$ /0.1% BSA and 1 μ l of DdeI (10 units, New England Biolabs) were added and the reannealed DNA was digested for 30 minutes at 56°C. One microliter of HinfI (10 units, New England Biolabs) was then added and incubated for another 30 minutes. The reaction was stopped by the addition of 4 μ l 75 mM EDTA and 6 μ l tracking dye to a final volume of 61 μ l.

[0125] The mineral oil was extracted with 0.2 ml chloroform, and 18 μ l of the reaction mixture (45 ng genomic DNA) was loaded onto a 30% polyacrylamide mini-gel (19:1, Rio Rad) in a Hoeffer SE200 apparatus. The gel was electrophoresed at approximately 300 volts for one hour until the bromphenol blue dye front migrated to 3.0 cm off-origin. The top 1.5 cm of the gel was removed and the remaining gel was exposed for four days with one intensification screen at -70°C.

Discussion of Photograph (Figure 9)

[0126] Each lane contains 45 ng of amplified genomic DNA. Lane A contains Molt4 DNA; Lane B, CH12; Lane C, SC-1; and Lane D, GM2064. Molt4 represents the genotype of a normal individual with two copies of the β^A gene per cell (AA), CH12 is a clinical sample from a sickle cell carrier with one β^A and one β^S gene per cell (AS), and SC-1 represents the genotype of a sickle cell individual with two copies of the β^S gene per cell (SS). GM2064, which contains no beta- or delta-globin sequences, is present as a negative control.

[0127] As seen in the photograph, the DdeI-cleaved, β^A -specific octamer is present only in those DNA's containing the β^A gene (Lanes A and B) and the HinfI-cleaved, β^S -specific trimer is present only in those DNA's containing the β^S gene (Lanes B and C). The presence of both trimer and octamer (Lane B) is diagnostic for a sickle cell carrier and is distinguishable from a normal individual (Lane A) with only octamer and a sickle cell afflicted individual (Lane C) with only trimer.

[0128] As a comparison, repeating the experiment described above using non-amplified genomic DNA revealed that the amplification increased the sensitivity of detection by at least 1000 fold.

EXAMPLE 6

[0129] This example illustrates direct detection of a totally unpurified single copy gene in whole human DNA on gels without the need for a labeled probe.

[0130] Using the technique described in Example 3, a 110-bp fragment from a sequence in the first exon of the beta-globin gene was amplified from 10 micrograms of whole human DNA after 20 cycles. This 110-bp fragment produced

after 20 cycles was easily visualized on gels stained with ethidium bromide.

[0131] The sequence was not amplified when it was first cut with the restriction enzyme DdeI unless, as in the beta-globin S allele, the sequence does not contain the restriction site recognized by the enzyme.

EXAMPLE 7

[0132]

A. A total of 100 fmoles pBR328 containing a 1.9 kb insert from the human beta-globin A allele, 50 nmoles each alpha-³²P-dNTP at 500 Ci/mole, and 1 nmole of each of the primers used in Example 3 were dissolved in a solution containing 100 µl 30 mM Tris-acetate at pH 7.9, 60 mM sodium acetate, 100 mM dithiothreitol, and 10 mM magnesium acetate. This solution was brought to 100°C for two minutes and cooled to 25°C for one minute. A total of 1 µl containing 4.5 units Klenow fragment of *E. coli* DNA polymerase I and 0.09 units inorganic pyrophosphatase was added to prevent the possible build-up of pyrophosphate in the reaction mixture, and the reaction was allowed to proceed for two minutes at 25°C, after which the cycle of heating, cooling, adding enzyme, and reacting was repeated nine times. Ten-µl aliquots were removed and added to 1 µl 600 mM EDTA after each synthesis cycle. Each was analyzed on a 14% polyacrylamide gel in 90 mM Tris-borate and 2.5 mM EDTA at pH 8.3 and 24 volts/cm for 2.5 hours. The completed gel was soaked for 20 minutes in the same buffer with the addition of 0.5 µg/ml ethidium bromide, washed with the original buffer, and photographed in UV light using a red filter.

The 110-bp fragment produced was excised from the gel under ultraviolet light and the incorporated ³²P counted by Cerenkov radiation. An attempt to fit the data to an equation of the form: pmoles/10 µl = 0.01 [(1+y)^N-y^N-1], where N represents the number of cycles and y the fractional yield per cycle, was optimal with y = 0.619. This indicates that a significant amplification is occurring.

B. The above experiment was repeated except that 100 nmoles of each dNTP was added to a 100 µl reaction, no radiolabel was employed, and aliquots were not removed at each cycle. After 10 cycles the reaction was terminated by boiling for two minutes and rehybridization was performed at 57°C for one hour. The sequence of the 110-bp product was confirmed by subjecting 8 µl aliquots to restriction analysis by addition of 1 µl bovine serum albumin (25 mg/ml) and 1 µl of the appropriate restriction enzyme (HinfI, MnII, MstII, NcoI) and by reaction at 37°C for 15 hours. PAGE was performed as described above.

EXAMPLE 8

[0133] This example illustrates the use of different primers to amplify various fragments of pBR328 and 322.

A. The experiment described in Example 7A was repeated except using the following primers: d(TTGCTTCT-GACACAAGTGTGTTCACTAGC) and d(GCCTCACCACCAACTTCATCCACGTTCCACC) to produce a 130-bp fragment of pBR328.

B. The experiment described in Example 7A was repeated except using the following primers: d(GGTT-GGCCAATCTACTCCCAGG) and d(TGGTCTCCTTAAACCTGTCTTG) to produce a 262-bp fragment of pBR328. The reaction time was 20 minutes per cycle.

C. The experiment described in Example 88 was repeated except that 100 fmoles of an MstII digest of pBR328 containing a 1.9 kb insert from the human beta-globin S allele was used as initial template. This plasmid was cleaved several times by MstII but not inside the sequence to be amplified. In addition, the primers employed were as follows:

d(GGTTGGCCAATCTACTCCCAGG)

and

d(TAACCTTGATACCAACCTGCCC)

to produce a 240-bp fragment.

D. The experiment described in Example 7B was repeated except that 100 fmoles of an NruI digest of pBR322 was used as template, 200 nmoles of each dNTP were used in the 100 µl reaction, and the primers were:

d(TAGGCGTATCACGAGGCCCT)

and

d(CTTCCCCATCGGTGATGTCG)

to produce a 500-bp fragment from pBR322. Reaction times were 20 minutes per cycle at 37°C. Final rehybridization was 15 hours at 57°C. Electrophoresis was on a 4% agarose gel.

EXAMPLE 9

[0134] This example illustrates the invention process wherein an in vitro mutation is introduced into the amplified segment.

A. A total of 100 fmoles of pBR322 linearized with NruI, 1 nmole each of the primers:

d(CGCATTAAAGCTTATCGATG)

and

d(TAGGCGTATCACGAGGCCCT)

designed to produce a 75-bp fragment, 100 nmole each dNTP, in 100 µl 40 mM Tris at pH 8, 20 mM in MgCl₂, 5 mM in dithiothreitol, and 5 mg/ml bovine serum albumin were combined. The mixture was brought to 100°C for one minute, cooled for 0.5 minutes in a water bath at 23°C, whereupon 4.5 units Klenow fragment and 0.09 units inorganic pyrophosphatase were added, and a reaction was allowed to proceed for three minutes. The cycle of heating, cooling, adding enzymes, and reacting was repeated nine times. The tenth reaction cycle was terminated by freezing and an 8-µl aliquot of the reaction mixture was applied to a 4% agarose gel visualized with ethidium bromide.

B. The experiment described in Example 9A was repeated except that the oligonucleotide primers employed were:

d(CGCATTAAAGCTTATCGATG)

and

d(AATTAATACGACTCACTATAGGGAGATAGGCGTATCACGAGGCCCT).

These primers are designed to produce a 101-bp fragment, 26 nucleotides of which (in the second listed primer) are not present in pBR322. These nucleotides represent the sequence of the T7 promoter, which was appended to the 75-bp sequence from pBR322 by using the primer with 20 complementary bases and a 26-base 5' extension. The procedure required less than two hours and produced two picomoles of the relatively pure 101-bp fragment from 100 fmoles of pBR322.

The T7 promoter can be used to initiate RNA transcription. T7 polymerase may be added to the 101-bp fragment to produce single-stranded RNA.

EP 0 502 589 B1

C. The experiment described in Example 8D was repeated except that the oligonucleotide primers employed were as follows:

5 d(TAGGCGTATCACGAGGCCCT)

and

10 d(CCAGCAAGACGTAGCCCAGC)

to produce a 1000-bp fragment from pBR322.

15 D. The experiment described in Example 9C was repeated except that the oligonucleotide primers employed were as follows:

d(TAGGCGTATCACGAGGCCCT)

20 and

d(AATTAATACGACTCACTATAGGGAGATAGGCGTATCACGAGGCCCT)

25 so as to produce a 1026-bp fragment, 26 nucleotides of which (in the second listed primer) are not present in pBR322 and represent the T7 promoter described above. The promoter has been inserted adjacent to a 1000-bp fragment from pBR322.

30 [0135] The results indicate that a primer which is not a perfect match to the template sequence but which is nonetheless able to hybridize sufficiently to be enzymatically extended produces a long product which contains the sequence of the primer rather than the corresponding sequence of the original template. The long product serves as a template for the second primer to introduce an in vitro mutation. In further cycles this mutation is amplified with an undiminished efficiency, because no further mispaired primings are required. In this case, a primer which carries a non-complementary extension on its 5' end was used to insert a new sequence in the product adjacent to the template sequence being copied.

EXAMPLE 10

40 [0136] This example illustrates employing nested sets of primers to decrease the background in the amplification of single copy genes.

[0137] Whole human DNA homozygous for the wild-type β -globin allele was subjected to twenty cycles of amplification as follows: A total of 10 μ g DNA, 200 picomoles each of the primers:

45 d(ACACAACGTGTTCAC TAGC)

and

50 d(CAACTTCATCCACGTT CACC)

55 and 100 nanomoles each dNTP in 100 μ l of 30 mM Tris-acetate pH 7.9, 60 mM sodium acetate, 10 mM dithiothreitol, and 10 mM magnesium acetate were heated to 100°C for one minute, cooled to 25°C for one minute, and treated with 2 units Klenow fragment for two minutes. The cycle of heating, cooling and adding Klenow was repeated 19 times. A ten- μ l aliquot was removed from the reaction mixture and subjected to a further ten cycles of amplification using each of the primers:

d(CAGACACCATGGTGCACCTGACTCCTG)

and

d(CCCCACAGGGCAGTAACGGCAGACTTCTCC),


which amplify a 58-bp fragment contained within the 110-bp fragment produced above. This final ten cycles of amplification was accomplished by diluting the 10-μl aliquot into 90 μl of the fresh Tris-acetate buffer described above containing 100 nanomoles each dNTP and 200 pmoles of each primer. Reaction conditions were as above. After ten cycles a 10-μl aliquot (corresponding to 100 nanograms of the original DNA) was applied to a 6% NuSieve (FMC Corp.) agarose gel and visualized using ethidium bromide.

[0138] Figure 10 illustrates this gel illuminated with UV light and photographed through a red filter as is known in the art. Lane 1 is molecular weight markers. Lane 2 is an aliquot of the reaction described above. Lane 3 is an aliquot of a reaction identical to that described above, except that the original wild-type DNA was cleaved with DdeI prior to amplification. Lane 4 is an aliquot of a reaction identical to that described above, except that human DNA homozygous for the sickle betaglobin allele was treated with DdeI prior to amplification (the sickle allele does not contain a DdeI site in the fragment being amplified here). Lane 5 is an aliquot of a reaction identical to that described above, except that salmon sperm DNA was substituted for human DNA. Lane 6 is an aliquot of a reaction identical to that described above, except that the aliquot was treated with DdeI after amplification (DdeI should convert the 58-bp wild-type product into 27-and 31-bp fragments). Lane 7 is an aliquot of the Lane 4 material treated with DdeI after amplification (the 58-bp sickle product contains no DdeI site).

[0139] Detection of a 58-bp fragment representative of a single-copy gene from one microgram of human DNA using only ethidium bromide staining of an agarose gel requires an amplification of about 500,000-fold. This was accomplished by using the two nested sets of oligonucleotide primers herein. The first set amplifies the 110-bp fragment and the inner nested set amplifies a sub-fragment of this product up to the level of convenient detection shown in Figure 10. This procedure of using primers amplifying a smaller -sequence contained within the sequence being amplified in the previous amplification process and contained in the extension products of the other primers allows one to distinguish the wild-type from the sickle allele at the beta-globin locus without resorting to either radioisotopic or non-radioisotopic probe hybridization methodology such as that of Conner et al., PNAS (USA), 80:278 (1983) and Leary et al., PNAS (USA), 80:4045 (1983).

EXAMPLE 11

[0140] The present process is expected to be useful in detecting, in a patient DNA sample, a specific sequence associated with an infectious disease such as, e.g., Chlamydia using a biotinylated hybridization probe spanning the desired amplified sequence and using the process described in U.S. 4,358,535, supra. The biotinylated hybridization probe may be prepared by intercalation and irradiation of a partially double-stranded DNA with a 4'-methylene substituted 4,5'-8-trimethylpsoralen attached to biotin via a spacer arm of the formula:



where Y is O, NH or N-CHO, x is a number from 1 to 4, and y is a number from 2 to 4. Detection of the biotinyl groups on the probe may be accomplished using a streptavidin-acid phosphatase complex commercially obtainable from Enzo Biochem Inc. using the detection procedures suggested by the manufacturer in its brochure. The hybridized probe is seen as a spot of precipitated stain due to the binding of the detection complex, and the subsequent reaction catalyzed by acid phosphatase, which produces a precipitable dye.

EXAMPLE 12

[0141] In this example, the process of Example 7 was basically used to amplify a 119 base pair fragment on the human β-hemoglobin gene using the primers:

5' -CTTCTGcagCAACTGTGTTCACTAGC-3' (GH18)
 5' -CACaAgCTTCATCCACGTTCAACC-3' (GH19)

5 where lower case letters denote mismatches from wild-type sequence to create restriction enzyme sites. The full scheme is shown in Table I. Table I illustrates a diagram of the primers GH18 and GH19 which are used for cloning and sequencing a 119-base pair fragment of the human β -globin gene and which are designed to contain internal restriction sites. The start codon ATG is underlined. GH18 is a 26-base oligonucleotide complementary to the negative strand and contains an internal PstI site. GH19 is a 23-base oligonucleotide complementary to the plus strand and contains an internal HindIII recognition sequence. Arrows indicate the direction of extension by DNA polymerase I. The boxed sequences indicate the restriction enzyme recognition sequences of each primer. These primers were selected by first screening the regions of the gene for homology to the PstI and HindIII restriction sites of bacteriophage M13. The primers were then prepared as described in previous examples.

Deel 1

8143

III DEPT
MAY 11 1964

Amplification and Cloning

[0142] After twenty cycles of amplification of 1 microgram of human genomic DNA isolated from the cell line Molt 4 as described in Example 2, 1/14th of the reaction product was hybridized to the labeled β -globin specific oligonucleotide probe, RS06, of the sequence 5'-CTGACTCCTGAGGAGAAGTCTGCCGTTACTGCCCTGTGGG-3' using the methods described above for oligomer restriction. Following solution hybridization, the reaction mixture was treated with DdeI under restriction digestion conditions as described above, to produce an 8-base pair oligonucleotide. The amount of this 8-base pair product is proportional to the amount of amplified product produced. The digestion products were resolved on a 30% polyacrylamide gel and visualized by autoradiography.

[0143] Analysis of the autoradiogram revealed that the amplification was comparable in efficiency to that of amplification with primers PC03 (5'-ACACAAGTGTTCAGTACG-3') and PC04 (5'-CCACTGCACCTACTTCAAC-3'), which are complementary to the negative and positive strands, respectively, of the wild-type β -globin.

[0144] The amplified product was ethanol precipitated to desalt and concentrate the sample, redissolved in a restriction buffer of 10 mM Tris pH 8, 10 mM $MgCl_2$, 1 mM DTT, 100 mM $NaCl$, and simultaneously digested with PstI and HindIII. After digestion the sample was desalted with a Centricon 10 concentrator and ligated overnight at 12°C with 0.3 micrograms of the PstI/HindIII digested vector M13mp10w, which is publicly available from Boehringer-Mannheim.

[0145] The entire ligation mixture was transformed into E. coli strain JM103, which is publicly available from BRL in Bethesda, MD. The procedure followed for preparing the transformed strain is described in Messing, J. (1981) Third Cleveland Symposium on Macromolecules: Recombinant DNA, ed. A. Walton, Elsevier, Amsterdam, 143-153.

[0146] The transformation mixture was plated onto x-gal media for screening via plaque hybridization with nylon filters. The filters were probed with a β -globin-specific oligonucleotide probe RS24 of the sequence 5'-CCCACAGGGCAGTAACGGCAGACTTCTCCTCAGGAGTCAG-3' to determine the number of β -globin inserts. The filters were then reprobed with the primer PC04 to determine the total number of inserts.

Plating and Screening

[0147] Table II summarizes the plating and plaque hybridization data. The filters were probed with the primer PC04 to determine the percentage of inserts resulting from amplification and cloning; 1206 clear plaques (90% of total number of clear plaques) hybridized to the primer. Fifteen plaques hybridized to the β -globin specific probe RS24. The percentage of β -globin positive plaques among the amplified primer-positive plaques is approximately 1%.

TABLE II

Plate No.	Blue Plaques	No Inserts*	Inserts**	β -Globin Inserts
1	28	25	246	1
2	29	18	222	2
3	11	26	180	0
4	24	20	192	5
5	22	27	185	5
6	39	21	181	3
TOTAL	158	132	1206	15

% of plaques containing amplified sequences which contain β -globin insert = $15/1206 \times 100 = 1.24\%$

% of total plaques which contain β -globin insert = $15/1496 \times 100 = \text{ca. } 1\%$

% of total plaques which contain amplified sequences = $1206/1496 \times 100 = 0.8\%$

* Clear plaques which do not hybridize to primer PC04

** Clear plaques which hybridize to primer PC04

Restriction Enzyme and Southern Blot Analysis

[0148] DNA from phage DNA miniprep preparation of three β -globin positive and two β -globin negative (but PC04 primer positive) plaques were analyzed by restriction enzyme analysis. MstII digestion of DNA from M13 clones containing the amplified β -globin fragment should generate a characteristic 283 base-pair fragment. Following MstII digestion, the three β -globin positive clones all produced the predicted 283 base pair fragment, while the two clones which were positive only with the primer produced larger fragments.

[0149] The gel from this analysis was transferred to a MSI nylon filter and hybridized with a radiolabeled nick-translated β -globin probe prepared by standard nick translation methods as described by Rigby et al., J. Mol. Biol. (1977), 113:237-51. The only bands which hybridized to the β -globin probe were the three β -globin positive clones. The two other clones had inserts which did not hybridize to the β -globin probe.

Sequence Analysis

[0150] Ten β -globin positive clones which were shown by restriction enzyme analysis to contain the β -globin insert were sequenced using the M13-dideoxy sequencing method. Of the ten clones, nine were identical to the β -globin wild-type sequence. The other clone was identical to the δ -globin gene which had been shown to be amplified to only a small degree by the β -globin primers.

[0151] In conclusion, the modified linker primers were nearly as efficient as the unmodified primers in amplifying the β -globin sequence. The primers were able to facilitate insertion of amplified DNA into cloning vectors. Due to the amplification of other segments of the genome, only 1% of the clones contained hemoglobin sequences.

[0152] Nine of the ten clones were found to be identical to the published β -globin sequence, showing that the technique amplifies genomic DNA with high fidelity. One clone was found to be identical with the published δ -globin sequence, confirming that the primers are specific for the β -globin gene despite their having significant sequence homology with δ -globin.

[0153] When cloning was carried out with a 267 base pair fragment of the β -globin gene, cloning was effective only when dimethylsulfoxide was present (10% by volume at 37°C) in the amplification procedure.

[0154] Restriction site-modified primers were also used to amplify and clone and partially sequence the human N-ras oncogene and to clone 240-base pair segments of the HLA DQ- α and DQ- β genes. All of these amplifications were carried out in the presence of 10% by volume dimethylsulfoxide at 37°C. The primers for amplifying HLA DQ- α and DQ- β genes were much more specific for their intended targets than were the β -globin and DR- β primers, which, rather than giving a discrete band on an ethidium bromide stained agarose gel, produced only a smear. In addition, the HLA DQ- α primers produced up to 20% of clones, with amplified inserts which contained the desired HLA target fragment, whereas 1% of the β -globin clones contained the target sequence. The HLA DQ- α and DQ- β gene cloning was only effective when the DMSO was present and the temperature was elevated.

EXAMPLE 13

[0155] This example illustrates the use of the process herein to prepare the TNF gene of 494 base pairs starting from two oligonucleotides of 74 base pairs each.

PRIMERS

[0156] The primers employed were prepared by the method described in Example 2 and are identified below, each being 74 mers.

(TN10) 5'-CCTCGTCTACTCCCAGGTCCTCTTCAAGGGCCAAGGCTGCCCCGACTATGTGCTCCTCA-
CCCACACCGTCAGCC-3'

(TN11) 5'-GGCAGGGGCTCTTGACGGCAGAGAGGAGTTGACCTTCTCCTGGTAGGAGATGGCGAAG-
CGGCTGACGGTGTGG-3'

(LL09) 5'-CCTGGCCAATGGCATGGATCTGAAAGATAACCAGCTGGTGGTGCCAGCAGATGGCCTGT-
ACCTCGTCTACTCCC-3'

(LL12) 5'-CTCCCTGATAGATGGGCTCATACCAGGGCTTGAGCTCAGCCCCCTCTGGGGTGTCTTTC-
GGGCAGGGGCTCTTG-3'

(TN08) 5'-TGTAGCAAACCATCAAGTTGAGGAGCAGCTCGAGTGGCTGAGCCAGCGGGCCAATGCCC-
TCCTGGCCAATGGCA-3'

(TN13) 5'-GATACTTGGGCAGATTGACCTCAGCGCTGAGTTGGTCACCCTTCTCCAGCTGGAAGACC-
CCTCCCTGATAGATG-3'

(LL07) 5'-CCTTAAGCTTATGCTCAGATCATCTTCTCAAACTCGAGTGACAAGCCTGTAGCCCATG-
TTGTAGCAAACCATC-3'

5 (TN14) 5'-GCTCGGATCCTTACAGGGCAATGACTCCAAAGTAGACCTGCCAGACTCGGCAAAGTCG-
AGATACTTGGGCAGA-3'

OVERALL PROCEDURE

10

[0157]

I. Ten cycles of the protocol indicated below were carried out using primers TN10 and TN11, which interact as shown in the diagram below, step (a).

15

II. A total of 2 µl of the reaction mixture from Part I above was added to the primers LL09 and LL12. The protocol described below was carried out for 15 cycles, so that the primers would interact with the product of Part I as shown in the diagram below, step (b).

20

III. A total of 2 µl of the reaction mixture from Part II above was added to the primers TN08 and TN13. The protocol described below was carried out for 15 cycles, so that the primers would interact with the product of Part II as shown in the diagram below, step (c).

25

IV. A total of 2 µl of the reaction mixture from Part III above was added to the primers LL07 and LL14. The protocol described below was carried out for 15 cycles, so that the primers would interact with the product of Part III as shown in the diagram below, step (d).

PROTOCOL

30

[0158] Each reaction contained 100 µl of:

2 mM of each of dATP, dCTP, dGTP and TTP

3 µM of each of the primers used at that step

35

1 x polymerase buffer, (30 mM Tris-acetate, 60 mM Na-acetate, 10 mM Mg-acetate, 2.5 mM dithiothreitol) Each cycle constituted:

40

1) 1 min. in boiling water

2) 1 min. cooling at room temperature

3) add 1 µl (5 units) of the Klenow fragment of DNA polymerase

45

4) allow the polymerization reaction to proceed for 2 min. For the next cycle start again at step 1.

50

55


DIAGRAM

[0159]

5

a)


10




15

b)

20




25




intermediate state

30



only the sequence between 5' of LL09 and 5' of LL12 will be full length. The strands that contain TN10 and TN11 have non-growing 5' ends. Thus...

35




40

c)


45




50



55





Deposit of Materials

[0160] The cell line SC-1 (CTCC #0082) was deposited on March 19, 1985 with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852 USA, with ATCC Accession No. CRL#8756. The deposit of SC-1 was made pursuant to a contract between the ATCC and the assignee of this patent application, Cetus Corporation. The contract with ATCC provides for permanent availability of the progeny of this cell line to the public on the issuance of the U.S. patent describing and identifying the deposit or the publications or upon the laying open to the public of any U.S. or foreign patent application, whichever comes first, and for availability of the progeny of this cell line to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 CFR §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638). The assignee of the present application has agreed that if the cell line on deposit should die or be lost or destroyed when cultivated under suitable conditions, it will be promptly replaced on notification with a viable culture of the same cell line.

[0161] In summary, the present invention is seen to provide a process for detecting sequences in nucleic acids by first amplifying one or more specific nucleic acid sequences using a chain reaction in which primer extension products are produced which can subsequently act as templates for further primer extension reactions. The process is especially useful in detecting nucleic acid sequences which are initially present in only very small amounts. Also, the amplification process can be used for molecular cloning.

Claims

1. An exponential amplification and detection kit for the amplification and detection of (a) specific template nucleic acid sequence(s) contained in a single- or double-stranded nucleic acid or in a mixture of such nucleic acids in a sample, which kit comprises in packaged form:

(a) at least a first and a second oligonucleotide primer different from each other, wherein

(aa) one of said primers is substantially complementary to said single-stranded nucleic acid or to one strand of said double-stranded nucleic acid,

(ab) the other primer of said primers is substantially complementary to a complement of said single-stranded nucleic acid or to the other strand of said double-stranded nucleic acid, and wherein

(ac) said primers define the termini of the specific nucleic acid sequence to be amplified and detected;

(b) an agent for polymerization; and

(c) means for detecting the amplified specific nucleic acid sequence.

2. The kit according to claim 1, wherein said specific template nucleic acid sequence is contained within a larger sequence.

3. The kit according to claim 1 or 2, wherein said nucleic acid is DNA or RNA, including messenger RNA, which DNA or RNA may be single-stranded or double-stranded, or is a DNA-RNA hybrid.
4. The kit according to any one of claims 1 to 3, wherein said nucleic acid is genomic DNA.
5. The kit according to any one of claims 1 to 4, wherein said primers contain about 15 to 25 nucleotides.
6. The kit according to any one of claims 1 to 5, wherein at least one of said primers contains a nucleotide sequence attached to its 5'-end which sequence is non-complementary to said nucleic acid.
7. The kit according to claim 6, wherein said nucleotide sequence attached to the 5'-end of said primer is a promoter sequence.
8. The kit of claim 7, wherein said promoter is the T7 promoter.
9. The kit according to any one of claims 1 to 5, wherein at least one of the primers is interspersed with bases or a nucleotide sequence which bases or nucleotide sequence are non-complementary to said specific nucleic acid sequence to be amplified and detected.
10. The kit according to any one of claims 1 to 5, wherein one primer of said primers contains a restriction site.
11. The kit according to any one of claims 1 to 10, wherein said agent for polymerization is an enzyme selected from E. coli DNA polymerase I, Klenow fragment of E. coli DNA polymerase I, T4 DNA polymerase, another DNA polymerase, reverse transcriptase, and heat-stable enzymes.
12. The kit according to any one of claim 1 to 11, wherein said means for detecting the amplified specific template nucleic acid sequence(s) comprises a labelled oligonucleotide probe.
13. Use of a kit according to any one of claims 1 to 12 for the amplification, detection and/or characterization of (a) specific template nucleic acid sequence(s) contained in a single- or double-stranded nucleic acid or in a mixture of such nucleic acids.
14. The use according to claim 13, wherein said specific template nucleic sequence(s) is (are) associated with infectious diseases such as those caused by bacteria, viruses and protozoan parasites, genetic disorders such as those caused by specific deletions and/or mutations in genomic DNA or cellular disorders such as cancer.

Patentansprüche

1. Exponentielle Amplifikation und Nachweis-Kit für die Amplifikation und den Nachweis in einer Probe von (einer) spezifischen Matrizen-Nucleinsäuresequenz(en), die in einer einzelsträngigen oder doppelsträngigen Nucleinsäure oder in einem Gemisch solcher Nucleinsäuren enthalten ist (sind), welcher in abgepackter Form folgende Bestandteile enthält:
 - (a) mindestens einen ersten und zweiten einzelsträngigen Oligonucleotidprimer, welche sich voneinander unterscheiden, wobei
 - (aa) einer dieser Primer im Wesentlichen komplementär zu der einzelsträngigen Nucleinsäure oder zu einem Strang der doppelsträngigen Nucleinsäure ist,
 - (ab) der andere Primer im Wesentlichen komplementär zu einem anderen Komplement der einzelsträngigen Nucleinsäure oder zum anderen Strang der doppelsträngigen Nucleinsäure ist, und wobei
 - (ac) diese Primer die Endpunkte der zu amplifizierenden und nachzuweisenden spezifischen Nucleinsäuresequenz definieren;
 - (b) ein Agens für die Polymerisation; und
 - (c) Mittel für den Nachweis der amplifizierten spezifischen Nucleinsäuresequenz.
2. Kit nach Anspruch 1, wobei die spezifische Matrizen-Nucleinsäuresequenz in einer größeren Sequenz enthalten

ist.

3. Kit nach einem der Ansprüche 1 oder 2, wobei die Nucleinsäure DNA oder RNA ist, einschließlich Messenger-RNA, wobei die DNA oder RNA als einzelsträngige oder doppelsträngige Nucleinsäure oder als DNA-RNA-Hybrid vorliegt.
4. Kit nach einem der Ansprüche 1 bis 3, wobei die Nucleinsäure genomische DNA ist.
5. Kit nach einem der Ansprüche 1 bis 4, wobei die Primer etwa 15 bis 25 Nucleotide umfassen.
6. Kit nach einem der Ansprüche 1 bis 5, wobei mindestens einer der Primer eine an seinem 5'-Ende angehängte Nucleotidsequenz enthält, deren Sequenz nicht komplementär zu der Nucleinsäure ist.
7. Kit nach Anspruch 6, wobei die am 5'-Ende des Primers angehängte Nucleotidsequenz eine Promotorsequenz ist.
8. Kit nach Anspruch 7, wobei der Promotor ein T7-Promotor ist.
9. Kit nach einem der Ansprüche 1 bis 5, wobei mindestens einer der Primer zusätzlich Basen oder eine Nucleotidsequenz enthält, wobei die Basen oder die Nucleotidsequenz zu der spezifischen Nucleinsäuresequenz, die amplifiziert und nachgewiesen werden soll, nicht komplementär sind.
10. Kit nach einem der Ansprüche 1 bis 5, wobei einer der Primer eine Restriktionsschnittstelle enthält.
11. Kit nach einem der Ansprüche 1 bis 10, wobei das Agens für die Polymerisation ein Enzym ist, ausgewählt aus E. coli-DNA-Polymerase I, Klenow-Fragment von E. coli-DNA-Polymerase I, T4-DNA-Polymerase, einer anderen DNA-Polymerase, reverser Transkriptase und hitzestabilen Enzymen.
12. Kit nach einem der Ansprüche 1 bis 11, wobei das Mittel für den Nachweis der amplifizierten spezifischen Matrizen-Nucleinsäuresequenz(en) eine markierte Oligonucleotidsonde umfasst.
13. Verwendung eines Kits nach einem der Ansprüche 1 bis 12, für die Amplifikation, den Nachweis und/oder die Charakterisierung von (einer) spezifischen Matrizen-Nucleinsäuresequenz(en), die in einer einzelsträngigen oder doppelsträngigen Nucleinsäure oder einem Gemisch solcher Nucleinsäuren enthalten ist (sind).
14. Verwendung nach Anspruch 13, wobei die spezifische(n) Matrizen-Nucleinsäuresequenz(en) mit infektiösen Erkrankungen, wie solche, die durch Bakterien, Viren und protozoische Parasiten verursacht werden, genetischen Störungen, wie solche, die durch spezifische Deletion und/oder Mutation in der genomischen DNA verursacht werden, oder zellulären Störungen, wie Krebs, in Verbindung stehen.


Revendications

1. Nécessaire d'amplification exponentielle et de détection pour l'amplification et la détection d'une(de) séquence(s) particulière(s) d'acide nucléique matrice contenue(s) dans un acide nucléique simple brin ou double brin ou dans un mélange de tels acides nucléiques dans un échantillon, lequel nécessaire comprend, sous forme conditionnée,
 - (a) au moins une première amorce oligonucléotidique et une seconde, différentes l'une de l'autre, dans lequel
 - (aa) l'une desdites amorces est pratiquement complémentaire dudit acide nucléique simple brin ou d'un brin dudit acide nucléique double brin,
 - (ab) l'autre amorce desdites amorces est pratiquement complémentaire d'un complément dudit acide nucléique simple brin ou de l'autre brin dudit acide nucléique double brin, et dans lequel
 - (ac) lesdites amorces définissent les extrémités de la séquence d'acide nucléique matrice particulière à amplifier et détecter ;
 - (b) un agent de polymérisation ; et
 - (c) des moyens pour détecter la séquence particulière d'acide nucléique amplifiée.

2. Nécessaire selon la revendication 1, dans lequel ladite séquence d'acide nucléique matrice particulière est contenue dans une séquence plus longue.
- 5 3. Nécessaire selon la revendication 1 ou 2, dans lequel ledit acide nucléique est un ADN ou un ARN, y compris un ARN messager, lequel ADN ou ARN peut être simple brin ou double brin, ou est un hybride ADN-ARN.
4. Nécessaire selon l'une quelconque des revendications 1 à 3, dans lequel ledit acide nucléique est un ADN génomique.
- 10 5. Nécessaire selon l'une quelconque des revendications 1 à 4, dans lequel lesdites amorces contiennent environ 15 à 25 nucléotides.
6. Nécessaire selon l'une quelconque des revendications 1 à 5, dans lequel au moins l'une desdites amorces contient une séquence nucléotidique attachée à son extrémité 5', laquelle séquence est non complémentaire dudit acide nucléique.
- 15 7. Nécessaire selon la revendication 6, dans lequel ladite séquence nucléotidique attachée à l'extrémité 5' de ladite amorce est une séquence de promoteur.
- 20 8. Nécessaire de la revendication 7, dans lequel ledit promoteur est le promoteur de T7.
9. Nécessaire selon l'une quelconque des revendications 1 à 5, dans lequel au moins l'une des amorces est parsemée de bases d'une séquence nucléotidique, lesquelles bases ou laquelle séquence nucléotidique sont non complémentaires de ladite séquence particulière d'acide nucléique à amplifier et à détecter.
- 25 10. Nécessaire selon l'une quelconque des revendications 1 à 5, dans lequel une amorce desdites amorces contient un site de restriction.
- 30 11. Nécessaire selon l'une quelconque des revendications 1 à 10, dans lequel ledit agent de polymérisation est une enzyme choisie parmi l'ADN polymérase I de *E. coli*, le fragment de Klenow d'ADN polymérase I de *E. coli*, l'ADN polymérase de T4, une autre ADN polymérase, la transcriptase inverse et des enzymes thermostables.
- 35 12. Nécessaire selon l'une quelconque des revendications 1 à 11, dans lequel lesdits moyens pour la détection de la (des) séquence(s) particulière(s) d'acide nucléique matrice amplifiée(s) comprennent une sonde oligonucléotidique marquée.
- 40 13. Utilisation d'un nécessaire selon l'une quelconque des revendications 1 à 12, pour l'amplification, la détection et/ou la caractérisation d'une (de) séquence(s) particulière(s) d'acide nucléique matrice contenue(s) dans un acide nucléique simple brin ou double brin ou dans un mélange de tels acides nucléiques.
- 45 14. Utilisation selon la revendication 13, dans laquelle ladite(lesdites) séquence(s) particulière(s) d'acide nucléique matrice particulière(s) est(sont) associée(s) à des maladies infectieuses, telles que celles provoquées par des bactéries, des virus ou des parasites de type protozoaire, à des troubles génétiques tels que ceux provoqués par des délétions et/ou des mutations particulières dans de l'ADN génomique, ou à des troubles cellulaires tels que le cancer.
- 50
- 55

FIG. 1

Double-Stranded 94-bp Sequence






FIG.2




FIG.3

Human Betaglobin

0 CCATCATATG CTTCATATG CTTCACAC AACTGTGTC ACTACCAAC ACCTGACTCC TCAAGCAAG TCTGCECTTA CTCCTCTCTG CGGCAAGCTG AACCTTCGCTG

101 111 121 131 *** 141 151 161 171 181 191 201 211 221

0 GSTAGCATAC GAATGTAAAC GAAGACTGTG TTGACACAAG TCATGCTGTG TGCACTAGAG ACTGCTCTTC AGACGCAAT GACGCGACAA CCGCTTCCAC TTCAACCAAC

PC01	PC02
CACAGGCAQTAAAG	TTTCTTCTGACACA

CYCLES

Signature, re-annual

CCATCTATTG CTTCATATG CTTCAGAAC AACTGTGTTC ACTAGCAACC ACTCTACTCC TGAGGAGAG TCTGCGCTTA CTGCGCTGTC GCGCAAGSTG AAGTTCGTCTG

extends <--- GCAAT GACCGGACAC 5' PCOI

Polymerase. dNTPs

0GCATCATTTG CTTCTACAC AACCTGTTTC ACTACCAACC ACCTCACTCC TGACGAAAG TCTCCCTTIA CTCCTCTCTG GCGCAAGGTG AACCTTGCTGTC

1GGTACATAAC GAATGTAAAC GAAGACTGTG TTACACAAG TGATCTGTGG TGGACTCAAG ACCTCTCTTC AGAGCGCAAT GACGGGACAC

1 TTTG CTCTGCAC AACCTGTTTC ACTACCAACC ACCTCACTCC TGACGAAAG TCTCCCTTIA CTCCTCTCTG GCGCAAGGTG AACCTTGCTGTC

0GGTACATAAC GAAGACTGTG TTACACAAG TGATCTGTGG TGGACTCAAG ACCTCTCTTC AGAGCGCAAT GACGGGACAC TTTCAACCAACC

denature, re-anneal

0 CCACTAATG CTTACATTTG CTCTGCACAC AACTCTGTC ACTACCAACC ACCTCACATC TCACGAAAC ICICGCTTA CTGCGCTGTC DCCCAAGCTC AACTTCCTCC

2

```

1      TTTC CTCTGCAC AACTCGTTT ACTAGCAACC ACCTCAGCTC TCGAGAGAAG TCATCCCGTA CTCCTCTGTG GGGCAAGCTG AAGTTGGTGQ
                                :       :       :       :       :       :       :       :       :
                                extends C----- CCAAT CAGCCGACAC          PCOI

```

Polymerase. DNTPs

[illegible]

FIG.4-3

Polymerase. dNTPs

0	CCAICTAATG	CTTCTGAC	AACTGTGTT	ACTAGCAAC	ACCTGACTCC	TCAGGAGAAG	TCGCGCGTTA	CTGCGCTGTG	GGCAGAAGTG	AACTTGCTGC
3	GCTAGATAAC	GAATGTAAAC	GAAGACTGTC	TTGACACAAG	TCATCTCTTC	TGCACTGAGG	ACTCCCTCTT	AGACGGCAAT	CACCGGACAC	
3	TTTG	CTTCTGAC	AACTGTGTT	ACTAGCAAC	ACCTGACTCC	TCAGGAGAAG	TCGCGCGTTA	CTGCGCTGTG		
2	GCTAGATAAC	GAATGTAAAC	GAAGACTGTC	TTGACACAAG	TCATCTCTTC	TGCACTGAGG	ACTCCCTCTT	AGACGGCAAT	CACCGGACAC	
2	TTTG	CTTCTGAC	AACTGTGTT	ACTAGCAAC	ACCTGACTCC	TCAGGAGAAG	TCGCGCGTTA	CTGCGCTGTG		
3	AAAC	GAAGACTGTC	TTGACACAAG	TCATCTCTTC	TGCACTGAGG	ACTCCCTCTT	AGACGGCAAT	CACCGGACAC		
3	TTTG	CTTCTGAC	AACTGTGTT	ACTAGCAAC	ACCTGACTCC	TCAGGAGAAG	TCGCGCGTTA	CTGCGCTGTG		
1	GCTAGATAAC	GAATGTAAAC	GAAGACTGTC	TTGACACAAG	TCATCTCTTC	TGCACTGAGG	ACTCCCTCTT	AGACGGCAAT	CACCGGACAC	
1	TTTG	CTTCTGAC	AACTGTGTT	ACTAGCAAC	ACCTGACTCC	TCAGGAGAAG	TCGCGCGTTA	CTGCGCTGTG	GGCAGAAGTG	AACTTGCTGC
3	AAAC	GAAGACTGTC	TTGACACAAG	TCATCTCTTC	TGCACTGAGG	ACTCCCTCTT	AGACGGCAAT	CACCGGACAC		
3	TTTG	CTTCTGAC	AACTGTGTT	ACTAGCAAC	ACCTGACTCC	TCAGGAGAAG	TCGCGCGTTA	CTGCGCTGTG		
2	AAAC	GAAGACTGTC	TTGACACAAG	TCATCTCTTC	TGCACTGAGG	ACTCCCTCTT	AGACGGCAAT	CACCGGACAC		
2	TTTG	CTTCTGAC	AACTGTGTT	ACTAGCAAC	ACCTGACTCC	TCAGGAGAAG	TCGCGCGTTA	CTGCGCTGTG	GGCAGAAGTG	AACTTGCTGC
3	AAAC	GAAGACTGTC	TTGACACAAG	TCATCTCTTC	TGCACTGAGG	ACTCCCTCTT	AGACGGCAAT	CACCGGACAC		
3	TTTG	CTTCTGAC	AACTGTGTT	ACTAGCAAC	ACCTGACTCC	TCAGGAGAAG	TCGCGCGTTA	CTGCGCTGTG	GGCAGAAGTG	AACTTGCTGC
0	GCTAGATAAC	GAATGTAAAC	GAAGACTGTC	TTGACACAAG	TCATCTCTTC	TGCACTGAGG	ACTCCCTCTT	AGACGGCAAT	CACCGGACAC	CCCTTCTCAC
										TTCAACACAC

COPIES OF DNA SEQUENCE AFTER N CYCLES

N	0	1	5	10	15	20
template	1	1	1	1	1	1
long product	0	1	5	10	15	20
short product	0	0	26	1013	32,752	1,048,555
-2(aspN)-N-1						




FIG.5

FIG.6

β^A	=====	CA TGG TGC ACC TGAC TCC TGAGGAGAAG TC TGCCG TTAC TGCCC TG TGGGGCAAGG TGAA GTACCACG TGGAC TGAGGAC TCC TC TTCAGACGGCAA TGACGGGACACCCCG TTCCAC TT	=====
β^S	===== *	CA TGG TGC ACC TGAC TCC TG TGGAGAAG TC TGCCG TTAC TGCCC TG TGGGGCAAGG TGAA GTACCACG TGGAC TGAGGACACC TC TTCAGACGGCAA TGACGGGACACCCCG TTCCAC TT	=====

* Marks the mutation (A to T) in the sickle cell gene which disrupts the DdeI site




FIG.7

* is label




FIG.9

A B C D



FIG. 10

